

LAMBERT SERIES OF LOGARITHM, THE DERIVATIVE OF DENINGER'S FUNCTION $R(z)$ AND A MEAN VALUE THEOREM FOR $\zeta\left(\frac{1}{2} - it\right)\zeta'\left(\frac{1}{2} + it\right)$

SOUMYARUP BANERJEE, ATUL DIXIT, AND SHIVAJEE GUPTA

Dedicated to Christopher Deninger on account of his 64th birthday

ABSTRACT. An explicit transformation for the series $\sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1}$, or equivalently, $\sum_{n=1}^{\infty} d(n) \log(n) e^{-ny}$ for $\operatorname{Re}(y) > 0$, which takes y to $1/y$, is obtained for the first time. This series transforms into a series containing the derivative of $R(z)$, a function studied by Christopher Deninger while obtaining an analogue of the famous Chowla-Selberg formula for real quadratic fields. In the course of obtaining the transformation, new important properties of $\psi_1(z)$ (the derivative of $R(z)$) are needed as is a new representation for the second derivative of the two-variable Mittag-Leffler function $E_{2,b}(z)$ evaluated at $b = 1$, all of which may seem quite unexpected at first glance. Our transformation readily gives the complete asymptotic expansion of $\sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1}$ as $y \rightarrow 0$ which was also not known before. An application of the latter is that it gives the asymptotic expansion of $\int_0^{\infty} \zeta\left(\frac{1}{2} - it\right)\zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt$ as $\delta \rightarrow 0$.

1. INTRODUCTION

Eisenstein series are the building blocks of modular forms and thus lie at the heart of the theory. In the case of the full modular group $\mathrm{SL}_2(\mathbb{Z})$, the Eisenstein series of even integral weight $k \geq 2$ are given by

$$E_k(z) := 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \frac{n^{k-1} q^n}{1 - q^n}, \quad (1.1)$$

where $q = e^{2\pi iz}$ with $z \in \mathbb{H}$ (the upper-half plane) and B_k are the Bernoulli numbers. They satisfy the modular transformations

$$\begin{aligned} E_k(z+1) &= E_k(z) \quad (k \geq 2), \\ E_k\left(\frac{-1}{z}\right) &= z^k E_k(z) \quad (k > 2), \quad E_2\left(\frac{-1}{z}\right) = z^2 E_2(z) + \frac{6z}{\pi i}. \end{aligned} \quad (1.2)$$

The series on the right-hand side of (1.1) is an example of what is known as a *Lambert series* whose general form is

$$\sum_{n=1}^{\infty} \frac{a(n)q^n}{1 - q^n} = \sum_{n=1}^{\infty} \frac{a(n)}{e^{ny} - 1} = \sum_{n=1}^{\infty} (1 * a)(n) e^{-ny}, \quad (1.3)$$

where $q = e^{-y}$ with $\operatorname{Re}(y) > 0$, and $a(n)$ is an arithmetic function with $(1 * a)(n) = \sum_{d|n} a(d)$ as the Dirichlet convolution.

For $\operatorname{Re}(s) > 1$, the Riemann zeta function is defined by $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$. Ramanujan [53, p. 173, Ch. 14, Entry 21(i)], [52, p. 319-320, formula (28)], [9, p. 275-276] derived a beautiful transformation involving the Lambert series associated with $a(n) = n^{-2m-1}$, $m \in \mathbb{Z} \setminus \{0\}$, and the odd zeta value

2020 *Mathematics Subject Classification*. Primary 11M06, 11N37; Secondary 33E12.
Keywords and phrases. Lambert series, Deninger's function, mean value theorems, asymptotic expansions.

$\zeta(2m+1)$, namely, for $\operatorname{Re}(\alpha), \operatorname{Re}(\beta) > 0$ with $\alpha\beta = \pi^2$,

$$\alpha^{-m} \left\{ \frac{1}{2} \zeta(2m+1) + \sum_{n=1}^{\infty} \frac{n^{-2m-1}}{e^{2\alpha n} - 1} \right\} = (-\beta)^{-m} \left\{ \frac{1}{2} \zeta(2m+1) + \sum_{n=1}^{\infty} \frac{n^{-2m-1}}{e^{2\beta n} - 1} \right\} - 2^{2m} \sum_{j=0}^{m+1} \frac{(-1)^j B_{2j} B_{2m+2-2j}}{(2j)!(2m+2-2j)!} \alpha^{m+1-j} \beta^j. \quad (1.4)$$

Along with the transformations of the Eisenstein series in (1.2), this formula also encapsulates the transformations of the corresponding Eichler integrals of the Eisenstein series as well as the transformation property of the Dedekind eta-function. The literature on this topic is vast with many generalizations and analogues for other L -functions, for example, [4], [6], [7], [12], [20], [21], [22], [23], [24], [32], [41], [42], and [44]. See also the recent survey article [10].

Recently, Kesarwani, Kumar and the second author [23, Theorem 2.4] obtained a new generalization of (1.4), namely, for $\operatorname{Re}(y) > 0$ and any complex a such that $\operatorname{Re}(a) > -1$,

$$\begin{aligned} & \sum_{n=1}^{\infty} \sigma_a(n) e^{-ny} + \frac{1}{2} \left(\left(\frac{2\pi}{y} \right)^{1+a} \operatorname{cosec} \left(\frac{\pi a}{2} \right) + 1 \right) \zeta(-a) - \frac{1}{y} \zeta(1-a) \\ &= \frac{2\pi}{y \sin \left(\frac{\pi a}{2} \right)} \sum_{n=1}^{\infty} \sigma_a(n) \left(\frac{(2\pi n)^{-a}}{\Gamma(1-a)} {}_1F_2 \left(1; \frac{1-a}{2}, 1 - \frac{a}{2}; \frac{4\pi^4 n^2}{y^2} \right) - \left(\frac{2\pi}{y} \right)^a \cosh \left(\frac{4\pi^2 n}{y} \right) \right), \end{aligned} \quad (1.5)$$

where ${}_1F_2(a; b, c; z)$ is the generalized hypergeometric function

$${}_1F_2(a; b, c; z) := \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n (c)_n} \frac{z^n}{n!},$$

where $z \in \mathbb{C}$ and $(a)_n = a(a+1) \cdots (a+n-1)$. They [23, Theorem 2.5] also analytically continued this result to $\operatorname{Re}(a) > -2m-3$, $m \in \mathbb{N} \cup \{0\}$, and in this way, they were able to get as corollaries not only Ramanujan's formula (1.4) and the transformation formula for the Dedekind eta-function but also new transformations when a is an even integer. Moreover, they showed [23, Equation (2.19)] that letting $a \rightarrow 0$ in (1.5) gives the following transformation of Wigert [62, p. 203, Equation (A)]:

$$\sum_{n=1}^{\infty} \frac{1}{e^{ny} - 1} = \frac{1}{4} + \frac{\gamma - \log(y)}{y} + \frac{2}{y} \sum_{n=1}^{\infty} \left\{ \log \left(\frac{2\pi n}{y} \right) - \frac{1}{2} \left(\psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) \right) \right\}, \quad (1.6)$$

where $\psi(z) := \Gamma'(z)/\Gamma(z)$ is the logarithmic derivative of the gamma function commonly known as the digamma function. Wigert [62, p. 203] called this transformation '*la formule importante*' (an important formula). Indeed, it is important, for, if we let $y \rightarrow 0$ in any angle $|\arg(y)| \leq \lambda$, where $\lambda < \pi/2$, it gives the complete asymptotic expansion upon using (3.13), that is,

$$\sum_{n=1}^{\infty} d(n) e^{-ny} \sim \frac{1}{4} + \frac{(\gamma - \log(y))}{y} - \sum_{n=1}^{\infty} \frac{B_{2n}^2 y^{2n-1}}{(2n)(2n)!}, \quad (1.7)$$

where γ is Euler's constant, which, in turn, readily implies [58, p. 163, Theorem 7.15]

$$\sum_{n=1}^{\infty} d(n) e^{-ny} = \frac{1}{4} + \frac{\gamma - \log y}{y} - \sum_{n=0}^{N-1} \frac{B_{2n+2}^2}{(2n+2)(2n+2)!} y^{2n+1} + O(|y|^{2N}). \quad (1.8)$$

The study of the moments

$$M_k(T) := \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt \quad (1.9)$$

is of fundamental importance in the theory of the Riemann zeta function. It is conjectured that $M_k(T) \sim C_k T \log^{k^2}(T)$ as $T \rightarrow \infty$ for positive constants C_k although for $k = 1$ and 2 , this has been proved by

Hardy and Littlewood [33] and Ingham [35] respectively. Such results are known as mean-value theorems for the zeta function. The importance of the study of moments lies, for example, in the fact that the estimate $M_k(T) = O_{k,\epsilon}(T^{1+\epsilon})$ for every natural number k is equivalent to the Lindelöf hypothesis [56] (see also [34]).

Another set of mean-value theorems which plays an important role in the theory is the one concerning the asymptotic behavior of the smoothly weighted moments, namely,

$$\int_0^\infty \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} e^{-\delta t} dt \quad (1.10)$$

as $\delta \rightarrow 0$. The relation between the two types of moments in (1.9) and (1.10) is given by a result [58, p. 159] which states that if $f(t) \geq 0$ for all t and for a given positive m ,

$$\int_0^\infty f(t) e^{-\delta t} dt \sim \frac{1}{\delta} \log^m \left(\frac{1}{\delta} \right)$$

as $\delta \rightarrow 0$, then

$$\int_0^T f(t) dt \sim T \log^m(T)$$

as $T \rightarrow \infty$. For an excellent survey on the moments, we refer the reader to [38].

The asymptotic expansion of $\sum_{n=1}^\infty d(n) e^{-ny}$ as $y \rightarrow 0$ in (1.8) allows us to obtain the asymptotic estimate for the smoothly weighted second moment, namely, as $\delta \rightarrow 0$, for every natural number N , we have [58, p. 164, Theorem 7.15 (A)]

$$\int_0^\infty \left| \zeta \left(\frac{1}{2} + it \right) \right|^2 e^{-\delta t} dt = \frac{\gamma - \log(2\pi\delta)}{2 \sin(\delta/2)} + \sum_{n=0}^N c_n \delta^n + O(\delta^{N+1}), \quad (1.11)$$

where the c_n are constants and the constant implied by the big-O depends on N . A simple proof of (1.11) was given by Atkinson [3]. In fact, (1.7) gives the complete asymptotic expansion for this moment.

The primary goal of this paper is to give a non-trivial application of (1.5). Note that the complex variable a in (1.5) enables differentiation of (1.5) with respect to a , which is not possible in (1.4) or other such known results. Indeed, it is an easy affair to check that differentiating the series on the left-hand side of (1.5) with respect to a and then letting $a = 0$ gives $\sum_{n=1}^\infty \frac{\log(n)}{e^{ny} - 1}$, which, in view of (1.3), satisfies

$$\sum_{n=1}^\infty \frac{\log(n)}{e^{ny} - 1} = \sum_{n=1}^\infty \log \left(\prod_{d|n} d \right) e^{-ny} = \frac{1}{2} \sum_{n=1}^\infty d(n) \log(n) e^{-ny}.$$

Here, in the last step, we used an elementary result $\prod_{d|n} d = n^{d(n)/2}$; see, for example, [2, Exercise 10, p. 47].

What is surprising though is, differentiating the right-hand side of (1.5) with respect to a and then letting $a = 0$ leads to an explicit and interesting series involving a well-known special function. This special function deserves a separate mention and hence after its brief introduction here, the literature on it is discussed in detail in Section 3.

In a beautiful paper [17], Deninger comprehensively studied the function $R : \mathbb{R}^+ \rightarrow \mathbb{R}$ uniquely defined by the difference equation

$$R(x+1) - R(x) = \log^2(x) \quad (R(1) = -\zeta''(0)).$$

The function $R(x)$ is an analogue of $\log(\Gamma(x))$ in view of the fact that the latter satisfies the difference equation $f(x+1) - f(x) = \log(x)$ with the initial condition $f(1) = 0$. As noted in [17, Remark (2.4)], R can be analytically continued to

$$\mathbb{D} := \mathbb{C} \setminus \{x \in \mathbb{R} \mid x \leq 0\}.$$

The special function which appears in our main theorem, that is, in Theorem 1.1, is $\psi_1(z)$, which is essentially the derivative of Deninger's function $R(z)$ (see (3.5) below). For $z \in \mathbb{D}$, it is given by

$$\psi_1(z) = -\gamma_1 - \frac{\log(z)}{z} - \sum_{n=1}^{\infty} \left(\frac{\log(n+z)}{n+z} - \frac{\log(n)}{n} \right), \quad (1.12)$$

where γ_1 is the first Stieltjes constant.

We are now ready to state the main result of our paper which transforms the Lambert series of logarithm into an infinite series consisting of $\psi_1(z)$.

Theorem 1.1. *Let $\psi_1(z)$ be given in (1.12). Then, for $\operatorname{Re}(y) > 0$,*

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1} &= -\frac{1}{4} \log(2\pi) + \frac{1}{2y} \log^2(y) - \frac{\gamma^2}{2y} + \frac{\pi^2}{12y} \\ &\quad - \frac{2}{y} (\gamma + \log(y)) \sum_{n=1}^{\infty} \left\{ \log \left(\frac{2\pi n}{y} \right) - \frac{1}{2} \left(\psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) \right) \right\} \\ &\quad + \frac{1}{y} \sum_{n=1}^{\infty} \left\{ \psi_1 \left(\frac{2\pi in}{y} \right) + \psi_1 \left(-\frac{2\pi in}{y} \right) - \frac{1}{2} \left(\log^2 \left(\frac{2\pi in}{y} \right) + \log^2 \left(-\frac{2\pi in}{y} \right) \right) + \frac{y}{4n} \right\}. \end{aligned} \quad (1.13)$$

Equivalently,

$$\begin{aligned} y \sum_{n=1}^{\infty} \frac{\gamma + \log(ny)}{e^{ny} - 1} - \frac{1}{4} y \log(y) + y \left(\frac{1}{4} \log(2\pi) - \frac{\gamma}{4} \right) + \frac{1}{2} \log^2(y) - \frac{\gamma^2}{2} - \frac{\pi^2}{12} \\ = \sum_{n=1}^{\infty} \left\{ \psi_1 \left(\frac{2\pi in}{y} \right) + \psi_1 \left(-\frac{2\pi in}{y} \right) - \frac{1}{2} \left(\log^2 \left(\frac{2\pi in}{y} \right) + \log^2 \left(-\frac{2\pi in}{y} \right) \right) + \frac{y}{4n} \right\}. \end{aligned} \quad (1.14)$$

Remark 1.1. *That the series*

$$\sum_{n=1}^{\infty} \left\{ \psi_1 \left(\frac{2\pi in}{y} \right) + \psi_1 \left(-\frac{2\pi in}{y} \right) - \frac{1}{2} \left(\log^2 \left(\frac{2\pi in}{y} \right) + \log^2 \left(-\frac{2\pi in}{y} \right) \right) + \frac{y}{4n} \right\}$$

converges absolutely is clear from (5.13) below.

The exact transformation in (1.13) is an analogue of Wigert's result (1.6). This is evident from the fact that

$$\log \left(\frac{2\pi n}{y} \right) - \frac{1}{2} \left(\psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) \right) = -\frac{1}{2} \left\{ \psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) - \left(\log \left(\frac{2\pi in}{y} \right) + \log \left(-\frac{2\pi in}{y} \right) \right) \right\},$$

which should be compared with the summand of the second series on the right-hand side of (1.13).

Thus, (1.13) allows us to transform $\sum_{n=1}^{\infty} d(n) \log(n) e^{-ny}$ into series having a constant times $1/y$ in the arguments of the functions in their summands. This "modular" behavior has an instant application: it gives the complete asymptotic expansion of $\sum_{n=1}^{\infty} d(n) \log(n) e^{-ny}$ as $y \rightarrow 0$, which is given in the following result.

Theorem 1.2. *Let A denote the Glaisher-Kinkelin constant defined by [27], [28], [43], [60, p. 461, Equation (A.7)]*

$$\log(A) := \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n k \log(k) - \left(\frac{n^2}{2} + \frac{n}{2} + \frac{1}{12} \right) \log(n) + \frac{n^2}{4} \right\}.$$

As $y \rightarrow 0$ in $|\arg(y)| < \pi/2$,

$$\sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1} \sim \frac{1}{2y} \log^2(y) + \frac{1}{y} \left(\frac{\pi^2}{12} - \frac{\gamma^2}{2} \right) - \frac{1}{4} \log(2\pi) + \frac{y}{12} \left(\log A - \frac{1}{12} \right)$$

$$+ \sum_{k=2}^{\infty} \frac{B_{2k} y^{2k-1}}{k} \left\{ \frac{B_{2k}}{2(2k)!} \left(\gamma - \sum_{j=1}^{2k-1} \frac{1}{j} + \log(2\pi) \right) + \frac{(-1)^k \zeta'(2k)}{(2\pi)^{2k}} \right\}. \quad (1.15)$$

As seen earlier, Wigert's result (1.8) is useful in getting the asymptotic estimate for the smoothly weighted second moment of the zeta function on the critical line given in (1.11). It is now natural to ask whether our result in Theorem 1.2 has an application in the theory of moments. Indeed, (1.15) implies the following result.

Theorem 1.3. *As $\delta \rightarrow 0$, $|\arg(\delta)| < \pi/2$,*

$$\int_0^{\infty} \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt = \frac{-\log^2(2\pi\delta) + \gamma^2 - \frac{\pi^2}{6}}{4 \sin\left(\frac{\delta}{2}\right)} + \sum_{k=0}^{2m-2} d_k \delta^k + O(\delta^{2m-1}), \quad (1.16)$$

where the d_k are effectively computable constants and the constant implied by the big- O depends on m .

In fact, one can obtain the complete asymptotic expansion of the left-hand side of (1.16) using (1.15).

Mean value theorems involving the derivatives of the Riemann zeta function have been well-studied. For example, Ingham [35] (see also Gonek [29, Equation (2)]¹) showed that

$$\int_0^T \zeta^{(\mu)}\left(\frac{1}{2} + it\right) \zeta^{(\nu)}\left(\frac{1}{2} - it\right) dt \sim \frac{(-1)^{\mu+\nu} T}{\mu + \nu + 1} \log^{\mu+\nu+1}(T)$$

as $T \rightarrow \infty$, where $\mu, \nu \in \mathbb{N} \cup \{0\}$. See also [59, p. 102]. In particular, for $\mu = 1$ and $\nu = 0$, we have

$$\int_0^T \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) dt \sim \frac{-T}{2} \log^2(T)$$

as $T \rightarrow \infty$.

The proof of Theorem 1.1 is quite involved and first requires establishing several new results that may seem quite unexpected at first glance. These results are important in themselves and may have applications in other areas. Hence this paper is organized as follows.

We collect frequently used results in the next section. In §3, we first prove Theorem 3.2 which gives an asymptotic expansion of $\psi_1(z)$ followed by Theorem 3.3, a Kloosterman-type result for $\psi_1(z)$. Theorem 3.4 is the highlight of this section and is an analogue of (2.2) established in [22, Theorem 2.2]. §4 is devoted to obtaining a new representation for the second derivative of the two-variable Mittag-Leffler function $E_{2,b}(z)$ at $b = 1$. In §5, we prove our main results, that is, Theorems 1.1, 1.2 and 1.3. Finally, we conclude the paper with some remarks and directions for future research.

2. PRELIMINARIES

Stirling's formula in a vertical strip $\alpha \leq \sigma \leq \beta$, $s = \sigma + it$ states that [16, p. 224]

$$|\Gamma(s)| = (2\pi)^{\frac{1}{2}} |t|^{\sigma - \frac{1}{2}} e^{-\frac{1}{2}\pi|t|} \left(1 + O\left(\frac{1}{|t|}\right) \right) \quad (2.1)$$

uniformly as $|t| \rightarrow \infty$.

We will also need the following result established in [22, Theorem 2.2] which is valid for $\operatorname{Re}(w) > 0$:

$$\sum_{n=1}^{\infty} \int_0^{\infty} \frac{t \cos(t)}{t^2 + n^2 w^2} dt = \frac{1}{2} \left\{ \log\left(\frac{w}{2\pi}\right) - \frac{1}{2} \left(\psi\left(\frac{iw}{2\pi}\right) + \psi\left(-\frac{iw}{2\pi}\right) \right) \right\}. \quad (2.2)$$

Watson's lemma is a very useful result in the asymptotic theory of Laplace integrals $\int_0^{\infty} e^{-zt} f(t) dt$.

This result typically holds for $|\arg(z)| < \pi/2$. However, with additional restrictions on f , Watson's lemma is known to hold for extended sectors. For the sake of completeness, we include it here in the form given in [57, p. 14, Theorem 2.2].

¹There is a slight typo in the asymptotic formula on the right-hand side of this equation in that $(-1)^{\mu+\nu}$ is missing.

Theorem 2.1. *Let f be analytic inside a sector $D : \alpha < \arg(t) < \beta$, where $\alpha < 0$ and $\beta > 0$. For each $\delta \in (0, \frac{1}{2}\beta - \frac{1}{2}\alpha)$, as $t \rightarrow 0$ in the sector $D_\delta : \alpha + \delta < \arg(t) < \beta - \delta$, we have*

$$f(t) \sim t^{\lambda-1} \sum_{n=0}^{\infty} a_n t^n,$$

where $\operatorname{Re}(\lambda) > 0$. Suppose there exists a real number σ such that $f(t) = O(e^{\sigma|t|})$ as $t \rightarrow \infty$ in D_δ . Then the integral

$$F_\lambda(z) := \int_0^\infty e^{-zt} f(t) dt \tag{2.3}$$

or its analytic continuation, has the asymptotic expansion

$$F_\lambda(z) \sim \sum_{n=0}^{\infty} a_n \frac{\Gamma(n + \lambda)}{z^{n+\lambda}}$$

as $|z| \rightarrow \infty$ in the sector

$$-\beta - \frac{\pi}{2} + \delta < \arg(z) < -\alpha + \frac{\pi}{2} - \delta.$$

The many-valued functions $t^{\lambda-1}$ and $z^{n+\lambda}$ have their principal values on the positive real axis and are defined by continuity elsewhere.

Actually we will be using an analogue of the above theorem where the integrand in (2.3) has a logarithmic factor.

We will be using Parseval's theorem given next. Let $\mathfrak{F}(s) = M[f; s]$ and $\mathfrak{G}(s) = M[g; s]$ denote the Mellin transforms of functions $f(x)$ and $g(x)$ respectively and let $c = \operatorname{Re}(s)$. If $M[f; 1 - c - it] \in L(-\infty, \infty)$ and $x^{c-1}g(x) \in L[0, \infty)$, then Parseval's formula [49, p. 83] is given by

$$\int_0^\infty f(x)g(x)dx = \frac{1}{2\pi i} \int_{(c)} \mathfrak{F}(1-s)\mathfrak{G}(s) ds, \tag{2.4}$$

where the vertical line $\operatorname{Re}(s) = c$ lies in the common strip of analyticity of the Mellin transforms $\mathfrak{F}(1-s)$ and $\mathfrak{G}(s)$, and, here and throughout the sequel, we employ the notation $\int_{(c)}$ to denote the line integral $\int_{c-i\infty}^{c+i\infty}$.

3. NEW RESULTS ON $\psi_1(z)$

In [19], Dilcher studied in detail the generalized gamma function $\Gamma_k(z)$ which relates to the Stieltjes constant $\gamma_k, k \geq 0$, defined by²

$$\gamma_k := \lim_{n \rightarrow \infty} \left(\sum_{j=1}^n \frac{\log^k(j)}{j} - \frac{\log^{k+1}(n)}{k+1} \right), \tag{3.1}$$

in a similar way as the Euler gamma function $\Gamma(z)$ relates to the Euler constant $\gamma = \gamma_0$. Using [19, Equation (2.1)] and [17, Equation (2.3.1)], we see that Dilcher's $\Gamma_1(z)$ is related to Deninger's $R(z)$ by³

$$\log(\Gamma_1(z)) = \frac{1}{2}(R(z) + \zeta''(0)). \tag{3.2}$$

As mentioned by Deninger in [17, Remark (2.4)], contrary to Euler's Γ , the function $\exp(R(x))$, or equivalently $\Gamma_1(x)$, where $x > 0$, cannot be meromorphically continued to the whole complex plane. But $\Gamma_1(z)$ is analytic in $z \in \mathbb{D}$. It is this $\exp(R(x))$ that Languasco and Righi [45] call as the *Ramanujan-Deninger gamma function*. They have also given a fast algorithm to compute it.

²Note that Deninger's definition of γ_1 in [17, p. 174] involves an extra factor of 2 which is not present in conventional definition of γ_1 , that is, in the $k = 1$ case of (3.1).

³By analytic continuation, Equation (2.3.1) from [17] is valid for $z \in \mathbb{D}$.

Dilcher also defined the generalized digamma function $\psi_k(z)$ as the logarithmic derivative of $\Gamma_k(z)$. His Proposition 10 from [19] implies that for $z \in \mathbb{D}$,

$$\psi_k(z) = -\gamma_k - \frac{\log^k(z)}{z} - \sum_{n=1}^{\infty} \left(\frac{\log^k(n+z)}{n+z} - \frac{\log^k(n)}{n} \right), \quad (3.3)$$

where $k \in \mathbb{N} \cup \{0\}$. Its special case $k = 1$ has already been given in (1.12). The function $\psi_k(z)$ occurs in Entry 22 of Chapter 8 in Ramanujan's second notebook, see [8]. It is also used by Ishibashi [36] to construct the k^{th} order Herglotz function which, in turn, plays an important role in his evaluation of the Laurent series coefficients of a zeta function associated to an indefinite quadratic form. As noted by Ishibashi and Kanemitsu [37, p. 78],

$$\psi_k(z) = \frac{1}{k+1} R'_{k+1}(z),$$

where $R_k(z)$ is defined by [17, p. 173]

$$R_k(z) = (-1)^{k+1} \left(\frac{\partial^k}{\partial s^k} \zeta(0, z) - \zeta^{(k)}(0) \right), \quad (3.4)$$

and $R_2(z) = R(z)$ of Deninger. Thus,

$$\psi_1(z) = \frac{1}{2} R'(z), \quad (3.5)$$

which is also implied by (3.2). The function $\psi_k(z)$ is related to the Laurent series coefficients $\gamma_k(z)$ of the Hurwitz zeta function $\zeta(s, z)$ known as the *generalized Stieltjes constants*. To see this, from [5, Theorem 1], note that if

$$\zeta(s, z) = \frac{1}{s-1} + \sum_{k=0}^{\infty} \frac{(-1)^k \gamma_k(z)}{k!} (s-1)^k, \quad (3.6)$$

then⁴

$$\gamma_k(z) = \lim_{n \rightarrow \infty} \left(\sum_{j=0}^n \frac{\log^k(j+z)}{j+z} - \frac{\log^{k+1}(n+z)}{k+1} \right) \quad (3.7)$$

so that $\gamma_k(1) = \gamma_k$. Then from (3.1), (3.3), (3.7) and the fact [19, Lemma 1] that

$$\lim_{n \rightarrow \infty} (\log^{k+1}(n+z) - \log^{k+1}(n)) = 0 \quad (z \in \mathbb{D}),$$

it is not difficult to see that

$$\psi_k(z) = -\gamma_k(z), \quad (3.8)$$

which was also shown by Shirasaka [55, p. 136]. Further properties and applications of $\psi_k(z)$ are derived in [18].

We thus see that the literature on $R(z)$, that is, $R_2(z)$, and, in general, on $R_k(z)$, is growing fast. In the words of Ishibashi [36, p. 61], “*Deninger proved several analytic properties of $R_2(x)$ in order to familiarize and assimilate it as one of the most commonly used number-theoretic special functions, . . .*”. Languasco and Righi [45] have also given a fast algorithm to compute $\psi_1(x)$, $x > 0$. The papers of Berndt [5], Blagouchine [11], Coffey [15], Chatterjee and Khurana [14] (and also the references therein) on the generalized Stieltjes constants, which, in view of (3.8), are nothing but $-\psi_k(z)$, are also vast sources of information on them. However, none of the studies prior to our current work is devoted to applications of these constants in obtaining a “modular” transformation for the Lambert series of logarithm and its application in the theory of the moments of the Riemann zeta function given in Theorems 1.1 and 1.3 respectively.

⁴It is to be noted that Berndt includes the factor $\frac{(-1)^k}{k!}$ in the definition of $\gamma_k(z)$ and does not have it in the summand of (3.6).

Our first result of this section gives the asymptotic expansion of $\psi_1(z)$ for $z \in \mathbb{D}$. To accomplish it, we require a generalization of Watson's lemma which allows for a logarithmic factor in the integrand. Such an expansion seems to have been first obtained by Jones [40, p. 439] (see also [63, Equations (4.14), (4.15)]). Though we will be using the same expansion, it is useful to rigorously derive it as a special case of a more general result due to Wong and Wyman [63, Theorem 4.1] given in the following theorem. We note in passing that Rieckstins [54] has also obtained asymptotic expansions of integrals involving logarithmic factors.

Theorem 3.1. *For $\gamma \in \mathbb{R}$, define a function*

$$F(z) := \int_0^{\infty e^{i\gamma}} f(t) e^{-zt} dt.$$

Assume that $F(z)$ exists for some $z = z_0$. If

(1) *for each integer $N \in \mathbb{N} \cup \{0\}$*

$$f(t) = \sum_{n=0}^N a_n t^{\lambda_n - 1} P_n(\log t) + o(t^{\lambda_N - 1} (\log t)^{m(N)}),$$

as $t \rightarrow 0$ along $\arg(t) = \gamma$.

(2) *$P_n(\omega)$ is a polynomial of degree $m = m(n)$.*

(3) *$\{\lambda_n\}$ is a sequence of complex numbers, with $\Re(\lambda_{n+1}) > \Re(\lambda_n)$, $\Re(\lambda_0) > 0$, for all n such that n and $n+1$ are in $\mathbb{N} \cup \{0\}$.*

(4) *$\{a_n\}$ is a sequence of complex numbers.*

Then as $z \rightarrow \infty$ in $S(\Delta)$

$$F(z) \sim \sum_{n=0}^N a_n P_n(D_n) [\Gamma(\lambda_n) z^{-\lambda_n}] + o(z^{-\lambda_N} (\log z)^{m(N)}),$$

where $S(\Delta) : |\arg(ze^{i\gamma})| \leq \frac{\pi}{2} - \Delta$, and D_n is the operator $D_n := \frac{d}{d\lambda_n}$. This result is uniform in the approach of $z \rightarrow \infty$ in $S(\Delta)$.

Theorem 3.2. *Fix any $\Delta > 0$. Then as $z \rightarrow \infty$ with $|\arg(z)| \leq \pi - \Delta$,*

$$\psi_1(z) \sim \frac{1}{2} \log^2(z) - \frac{1}{2z} \log z + \sum_{k=1}^{\infty} \frac{B_{2k}}{2k z^{2k}} \left(\sum_{j=1}^{2k-2} \frac{1}{j} + \frac{1}{2k-1} - \log z \right). \quad (3.9)$$

Proof. We first prove the result for $|\arg(z)| < \pi/2$ and then extend it to $|\arg(z)| \leq \pi - \Delta$. To that end, we begin with the analogue for $R(z)$ of Plana's integral for $\log(\Gamma(z))$, namely, for $\Re(z) > 0$, we have [17, Equation (2.12)]

$$R(z) = -\zeta''(0) - 2 \int_0^{\infty} \left((z-1)e^{-t} + \frac{e^{-zt} - e^{-t}}{1 - e^{-t}} \right) \frac{\gamma + \log(t)}{t} dt. \quad (3.10)$$

Differentiating (3.10) under the integral sign with respect to z and using (3.5), we see that

$$\begin{aligned} \psi_1(z) &= - \int_0^{\infty} \left(e^{-t} - \frac{te^{-zt}}{1 - e^{-t}} \right) \frac{(\gamma + \log(t))}{t} dt \\ &= - \int_0^{\infty} (e^{-t} - e^{-zt}) (\gamma + \log(t)) \frac{dt}{t} - \int_0^{\infty} e^{-zt} \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) (\gamma + \log(t)) dt \\ &= \frac{1}{2} \log^2(z) - \int_0^{\infty} e^{-zt} \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) (\gamma + \log(t)) dt, \end{aligned} \quad (3.11)$$

where we used the fact that for $\operatorname{Re}(z) > 0$,

$$\int_0^\infty (e^{-t} - e^{-zt})(\gamma + \log(t)) \frac{dt}{t} = -\frac{1}{2} \log^2(z),$$

which follows from [17, Equation (2.13)]⁵

$$\int_0^\infty (e^{-\beta t} - e^{-\alpha t})(\gamma + \log(t)) \frac{dt}{t} = \frac{1}{2} (\log^2(\beta) - \log^2(\alpha)).$$

Thus

$$\psi_1(z) = \frac{1}{2} \log^2(z) - \gamma(\psi(z) - \log(z)) - \int_0^\infty e^{-zt} \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) \log(t) dt, \quad (3.12)$$

where we employed the well-known result [31, p. 903, Formula **8.361.8**] that for $\operatorname{Re}(z) > 0$,

$$\psi(z) = \log(z) + \int_0^\infty e^{-zt} \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) dt.$$

We now find the asymptotic expansion of the integral on the right-hand side of (3.12), that is, of

$$I := \int_0^\infty e^{-zt} f(t) dt,$$

where

$$f(t) := \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) \log(t),$$

by applying Theorem 3.1. To that end, observe that for $|t| < 2\pi$,

$$\begin{aligned} f(t) &= \log(t) \left(\frac{1}{t} - \frac{1}{1 - e^{-t}} \right) \\ &= -\frac{\log(t)}{t} \left(\frac{te^t}{e^t - 1} - 1 \right) \\ &= -\frac{\log(t)}{t} \left(\sum_{n=0}^\infty \frac{B_n(1)t^n}{n!} - 1 \right) \\ &= \log(t) \sum_{n=0}^\infty \frac{(-1)^n B_{n+1} t^n}{(n+1)!}, \end{aligned}$$

where $B_n(x)$ are Bernoulli polynomials. Thus, with $P_n(x) = x$, $\lambda_n = n + 1$, $a_n = \frac{(-1)^n B_{n+1}}{(n+1)!}$, $n \geq 0$, all of the hypotheses of Theorem 3.1 are satisfied, and hence

$$\begin{aligned} I &\sim \sum_{n=0}^\infty \frac{(-1)^n B_{n+1}}{(n+1)z^{n+1}} (\psi(n+1) - \log(z)) \\ &= \sum_{n=1}^\infty \frac{(-1)^{n-1} B_n}{nz^n} \left(-\gamma + \sum_{k=1}^{n-1} \frac{1}{k} - \log(z) \right), \end{aligned}$$

where we used the elementary fact $\psi(n) = -\gamma + \sum_{k=1}^{n-1} \frac{1}{k}$. Inserting this asymptotic expansion of I in (3.12) along with that of $\psi(z)$, namely, for $|\arg z| < \pi$,

$$\psi(z) = \log(z) - \frac{1}{2z} - \sum_{n=1}^\infty \frac{B_{2n}}{2nz^{2n}}, \quad (3.13)$$

⁵Deninger requires it with $\alpha > 0, \beta > 0$, however, it is easily seen to hold for $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\beta) > 0$ as well.

as $z \rightarrow \infty$, we arrive at

$$\begin{aligned}\psi_1(z) &\sim \frac{1}{2} \log^2(z) - \gamma \left(-\frac{1}{2z} - \sum_{n=1}^{\infty} \frac{B_{2n}}{2nz^{2n}} \right) + \sum_{n=1}^{\infty} \frac{(-1)^n B_n}{nz^n} \left(-\gamma + \sum_{j=1}^{n-1} \frac{1}{j} - \log(z) \right) \\ &= \frac{1}{2} \log^2(z) + \sum_{n=1}^{\infty} \frac{(-1)^n B_n}{nz^n} \left(\sum_{j=1}^{n-1} \frac{1}{j} - \log(z) \right) \\ &= \frac{1}{2} \log^2(z) - \frac{1}{2z} \log z + \sum_{n=1}^{\infty} \frac{B_{2n}}{2nz^{2n}} \left(\sum_{j=1}^{2n-2} \frac{1}{j} + \frac{1}{2n-1} - \log(z) \right)\end{aligned}$$

using the well-known facts $B_1 = -1/2$ and $B_{2n-1} = 0, n > 1$. This proves (3.9) for $|\arg(z)| < \pi/2$.

To extend it to $|\arg(z)| \leq \pi - \Delta$, where $\Delta > 0$, we use the analogue of Theorem 2.1 containing a logarithmic factor in the integrand of the concerned integral, which practically changes none of the hypotheses in the statement of Theorem 2.1 and its proof⁶ since $\log(t) = O(t^\epsilon)$ as $t \rightarrow \infty$ for any $\epsilon > 0$. We apply it with $\alpha = -\pi/2$ and $\beta = \pi/2$. It shows that the expansion in (3.9) holds for $|\arg(z)| \leq \pi - \Delta$. \square

Remark 3.1. *The result in the above theorem is not new. In fact, in [15, Proposition 3], this has been done for $\psi_k(z)$ for all $k \in \mathbb{N}$ in view of (3.8). We derive the asymptotic expansion for $\psi_1(z)$ right from scratch only to make this paper self-contained. Also, it is to be noted that in [15], the result has been proved only for $|\arg(z)| < \pi/2$ whereas we crucially require the result to hold for $|\arg(z)| \leq \pi - \Delta$ for $\Delta > 0$. We also note that one could also obtain this asymptotic behavior beginning with the Euler-Maclaurin summation formula.*

Our next result is a new analogue of Kloosterman's result for $\psi(x)$ [58, p. 24-25].

Theorem 3.3. *Let $|\arg(z)| < \pi$. Let ψ_1 be defined in (1.12). For $0 < c = \operatorname{Re}(s) < 1$,*

$$\psi_1(z+1) - \frac{1}{2} \log^2(z) = \frac{1}{2\pi i} \int_{(c)} \frac{\pi \zeta(1-s)}{\sin(\pi s)} (\gamma - \log(z) + \psi(s)) z^{-s} ds. \quad (3.14)$$

Proof. We first prove the result for $z > 0$ and later extend it to $|\arg(z)| < \pi$ by analytic continuation. Using (3.11), we have

$$\begin{aligned}\psi_1(z) - \frac{1}{2} \log^2(z) &= \int_0^\infty e^{-zt} (\gamma + \log(t)) dt + \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) e^{-zt} (\gamma + \log(t)) dt \\ &= -\frac{\log(z)}{z} + \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) e^{-zt} (\gamma + \log(t)) dt,\end{aligned} \quad (3.15)$$

where in the last step we employed [31, p. 573, formula **4.352.1**]

$$\int_0^\infty t^{s-1} e^{-zt} (\gamma + \log(t)) dt = \frac{\Gamma(s)}{z^s} (\gamma - \log(z) + \psi(s)) \quad (\operatorname{Re}(s) > 0) \quad (3.16)$$

with $s = 1$ and the fact that $\psi(1) = -\gamma$. We now evaluate the integral on the right-hand side of (3.15) by means of Parseval's formula for Mellin transforms [49, p. 83, Equation (3.1.13)]. For $0 < \operatorname{Re}(s) < 1$, we have [58, p. 23, Equation (2.7.1)]

$$\int_0^\infty t^{s-1} \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) dt = \Gamma(s) \zeta(s). \quad (3.17)$$

⁶Temme [57, p. 15] refers to Olver [47, p. 114] for a proof.

Therefore, along with (3.16) and the equation given above, for $0 < c = \operatorname{Re}(s) < 1$, Parseval's formula (2.4) implies

$$\begin{aligned} \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) e^{-zt} (\gamma + \log(t)) dt &= \frac{1}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) \frac{\Gamma(1-s)}{z^{1-s}} (\gamma - \log(z) + \psi(1-s)) ds \\ &= \frac{1}{2\pi i z} \int_{(c)} \frac{\pi \zeta(s)}{\sin(\pi s)} (\gamma - \log(z) + \psi(1-s)) z^s ds \\ &= \frac{1}{2\pi i} \int_{(c')} \frac{\pi \zeta(1-s)}{\sin(\pi s)} (\gamma - \log(z) + \psi(s)) z^{-s} ds, \end{aligned} \quad (3.18)$$

where $0 < c' < 1$. In the second step, we used the reflection formula $\Gamma(s)\Gamma(1-s) = \pi/\sin(\pi s)$, $s \notin \mathbb{Z}$, and in the last step, we replaced s by $1-s$.

Now (3.14) follows by substituting (3.18) in (3.15) and using the fact [19, Equation (8.3)]

$$\psi_1(z) = \psi_1(z+1) - \frac{\log(z)}{z}. \quad (3.19)$$

This proves (3.14) for $z > 0$. The result is easily seen to be true for any complex z such that $|\arg(z)| < \pi$ by analytic continuation with the help of (1.12), elementary bounds on the Riemann zeta function, Stirling's formula (2.1) and the corresponding estimate for $\psi(s)$. \square

In the next theorem, we give a closed-form evaluation of an infinite series of integrals in terms of the digamma function and $\psi_1(z)$. This theorem is an analogue of (2.2) and will play a fundamental role in the proof of Theorem 1.1.

Theorem 3.4. *For $\operatorname{Re}(w) > 0$, we have*

$$\begin{aligned} 4 \sum_{m=1}^{\infty} \int_0^\infty \frac{u \cos(u) \log(u/w)}{u^2 + (2\pi m w)^2} du &= \psi_1(iw) - \frac{1}{2} \log^2(iw) + \psi_1(-iw) - \frac{1}{2} \log^2(-iw) + \frac{\pi}{2w} \\ &\quad + \gamma (\psi(iw) + \psi(-iw) - 2 \log(w)). \end{aligned} \quad (3.20)$$

Proof. Using Theorem 3.3, once with $x = iw$, and again with $x = -iw$ and adding the respective sides, for $0 < c = \operatorname{Re}(s) < 1$, we obtain

$$\begin{aligned} &\psi_1(iw+1) - \frac{1}{2} \log^2(iw) + \psi_1(-iw+1) - \frac{1}{2} \log^2(-iw) \\ &= \frac{1}{2\pi i} \int_{(c)} \frac{\pi \zeta(1-s)}{\sin(\pi s)} (\gamma - \log(iw) + \psi(s)) (iw)^{-s} ds + \frac{1}{2\pi i} \int_{(c)} \frac{\pi \zeta(1-s)}{\sin(\pi s)} (\gamma - \log(-iw) + \psi(s)) (-iw)^{-s} ds \\ &= \frac{1}{2\pi i} \int_{(c)} \frac{2\pi \zeta(1-s)}{\sin(\pi s)} (\gamma + \psi(s)) \cos\left(\frac{\pi s}{2}\right) w^{-s} ds - I_2, \end{aligned} \quad (3.21)$$

where

$$I_2 := \frac{1}{2\pi i} \int_{(c)} \frac{\pi \zeta(1-s)}{\sin(\pi s)} \left(e^{-\frac{i\pi s}{2}} \log(iw) + e^{\frac{i\pi s}{2}} \log(-iw) \right) w^{-s} ds. \quad (3.22)$$

Next, using the series expansions of the exponential functions $e^{\frac{i\pi s}{2}}$ and $e^{-\frac{i\pi s}{2}}$ and splitting them according to n even and n odd, it is easily seen that

$$e^{-\frac{i\pi s}{2}} \log(iw) + e^{\frac{i\pi s}{2}} \log(-iw) = 2 \log(w) \cos\left(\frac{\pi s}{2}\right) + \pi \sin\left(\frac{\pi s}{2}\right). \quad (3.23)$$

Hence from (3.21), (3.22) and (3.23), we see that

$$\psi_1(iw+1) - \frac{1}{2} \log^2(iw) + \psi_1(-iw+1) - \frac{1}{2} \log^2(-iw) =: J_1 - J_2, \quad (3.24)$$

where

$$J_1 := \frac{1}{2\pi i} \int_{(c)} \frac{\pi \zeta(1-s)}{\sin\left(\frac{\pi s}{2}\right)} (\gamma + \psi(s) - \log(w)) w^{-s} ds,$$

$$J_2 := \frac{1}{2\pi i} \int_{(c)} \frac{\pi^2 \zeta(1-s)}{2 \cos\left(\frac{\pi s}{2}\right)} w^{-s} ds.$$

Using the functional equation of $\zeta(s)$ [58, p. 13, Equation (2.1.1)]

$$\zeta(s) = 2^s \pi^{s-1} \Gamma(1-s) \zeta(1-s) \sin\left(\frac{1}{2}\pi s\right), \quad (3.25)$$

with s replaced by $1-s$, we have

$$J_2 = \frac{\pi^2}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) (2\pi w)^{-s} ds = \pi^2 \left(\frac{1}{e^{2\pi w} - 1} - \frac{1}{2\pi w} \right), \quad (3.26)$$

where the last step follows from (3.17). Again using (3.25) with s replaced by $1-s$, we observe that

$$J_1 = \frac{1}{2\pi i} \int_{(c)} 2\pi \Gamma(s) \zeta(s) \cot\left(\frac{\pi s}{2}\right) (\gamma + \psi(s) - \log(w)) (2\pi w)^{-s} ds.$$

Now it is important to observe that shifting the line of integration from $\operatorname{Re}(s) = c$, $0 < c < 1$ to $\operatorname{Re}(s) = d$, $1 < d < 2$ does not introduce any pole of the integrand. So consider the rectangular contour $[c - iT, d - iT], [d - iT, d + iT], [d + iT, c + iT]$ and $[c + iT, c - iT]$. By Cauchy's residue theorem and the fact that the integral along the horizontal segments of the contour tend to zero as $T \rightarrow \infty$, as can be seen from (2.1), elementary bounds of the zeta function and the fact [29, Equation (15)] $\psi(s) = \log(s) + O(1/|s|)$, it is seen that

$$\begin{aligned} J_1 &= \frac{1}{2\pi i} \int_{(d)} 2\pi \Gamma(s) \zeta(s) \cot\left(\frac{\pi s}{2}\right) (\gamma + \psi(s) - \log(w)) (2\pi w)^{-s} ds \\ &= \sum_{m=1}^{\infty} \frac{1}{2\pi i} \int_{(d)} 2\pi \Gamma(s) \cot\left(\frac{\pi s}{2}\right) (\gamma + \psi(s) - \log(w)) (2\pi m w)^{-s} ds \\ &= 2\pi \sum_{m=1}^{\infty} \left\{ \frac{(\gamma - \log(w))}{2\pi i} \int_{(d)} \Gamma(s) \cot\left(\frac{\pi s}{2}\right) (2\pi m w)^{-s} ds + \frac{1}{2\pi i} \int_{(d)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) (2\pi m w)^{-s} ds \right\}, \end{aligned} \quad (3.27)$$

where in the last step, we used the series representation for $\zeta(s)$ and then interchanged the order of summation and integration which is valid because of absolute and uniform convergence. We now find convenient representations for the two line integrals.

For $0 < \operatorname{Re}(s) = (c_1) < 2$ and $\operatorname{Re}(z) > 0$, we show that

$$\frac{1}{2\pi i} \int_{(c_1)} \Gamma(s) \cot\left(\frac{\pi s}{2}\right) z^{-s} ds = \frac{2}{\pi} \int_0^{\infty} \frac{u \cos(u)}{u^2 + z^2} du. \quad (3.28)$$

We first prove this result for $z > 0$ and then extend it by analytic continuation. We also first prove the result for $0 < \operatorname{Re}(s) = c_1 < 1$. To that end, using the double angle formulas for sine and cosine, we have

$$\begin{aligned} \frac{1}{2\pi i} \int_{(c_1)} \Gamma(s) \cot\left(\frac{\pi s}{2}\right) z^{-s} ds &= \frac{1}{2\pi i} \int_{(c_1)} \Gamma(s) \frac{(1 + \cos(\pi s))}{\sin(\pi s)} z^{-s} ds \\ &= \frac{1}{2\pi i} \int_{(c_1)} \Gamma(s) \operatorname{cosec}(\pi s) z^{-s} ds + \frac{1}{2\pi i} \int_{(c_1)} \Gamma(s) \cot(\pi s) z^{-s} ds \\ &= -\frac{1}{\pi} e^z \operatorname{Ei}(-z) - \frac{1}{\pi} e^{-z} \operatorname{Ei}(z), \end{aligned} \quad (3.29)$$

where in the last step we used [13, p. 102, Formulas **3.3.2.1**], and where $\text{Ei}(x)$ is the exponential integral defined [51, p. 788] for $z > 0$ by $\text{Ei}(z) := \int_{-\infty}^z e^t/t dt$, or, as is seen from [39, p. 1], by $\text{Ei}(-z) := -\int_z^{\infty} e^{-t}/t dt$. Equation (3.28) now follows upon using [50, p. 395, Formula **2.5.9.12**]

$$e^{-z}\text{Ei}(z) + e^z\text{Ei}(-z) = -2 \int_0^{\infty} \frac{u \cos(u)}{u^2 + z^2} du. \quad (3.30)$$

Since both sides are analytic for $\text{Re}(z) > 0$, the result now follows by analytic continuation for $\text{Re}(z) > 0$.

Next, observe that shifting the line of integration from $\text{Re}(s) = c_1$ to $\text{Re}(s) = d$, where $1 < d < 2$, and using Cauchy's residue theorem does not change the evaluation of the integral on the left-hand side of (3.28). Hence invoking (3.28) with $z = 2\pi mw$, we find that

$$\begin{aligned} \sum_{m=1}^{\infty} \frac{1}{2\pi i} \int_{(d)} \Gamma(s) \cot\left(\frac{\pi s}{2}\right) (2\pi mw)^{-s} ds &= \frac{2}{\pi} \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u)}{u^2 + (2\pi mw)^2} du \\ &= \frac{1}{\pi} \left(\log(w) - \frac{1}{2} (\psi(iw) + \psi(-iw)) \right), \end{aligned} \quad (3.31)$$

where in the last step, we employed (2.2).

Here, it is important to note that the series representation of J_1 in (3.27) and the convergence of the series in (3.31) together imply that the series

$$\sum_{m=1}^{\infty} \frac{1}{2\pi i} \int_{(d)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) (2\pi mw)^{-s} ds$$

converges too. We now suitably transform the line integral in the above equation into an integral of a real variable by proving that for $0 < \text{Re}(s) = c < 1$ and $\text{Re}(z) > 0$, we have

$$\frac{1}{2\pi i} \int_{(c)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) z^{-s} ds = \frac{2}{\pi} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + z^2} du + \frac{\pi}{2} e^{-z}. \quad (3.32)$$

Again, we first prove this result for $z > 0$ and later extend it by analytic continuation to $\text{Re}(z) > 0$. Note that from [46, p. 201, Equation (5.68)], for $0 < \text{Re}(s) = c < 1$ and $x > 0$,

$$\frac{1}{2\pi i} \int_{(c)} \Gamma(s) \psi(s) \cos\left(\frac{\pi s}{2}\right) (xz)^{-s} ds = \cos(xz) \log(xz) + \frac{\pi}{2} \sin(xz). \quad (3.33)$$

Multiply both sides by $\frac{x}{1+x^2}$ and integrate with respect to x from 0 to ∞ thereby obtaining

$$\int_0^{\infty} \frac{x}{1+x^2} \frac{1}{2\pi i} \int_{(c)} \Gamma(s) \psi(s) \cos\left(\frac{\pi s}{2}\right) (xz)^{-s} ds dx = \int_0^{\infty} \left(\cos(xz) \log(xz) + \frac{\pi}{2} \sin(xz) \right) \frac{x dx}{1+x^2} \quad (3.34)$$

On the left-hand side, an application of Fubini's theorem allows us to interchange the order of integration. On the right-hand side, we employ the change of variable $x = u/z$. Doing this and then using the well-known evaluation

$$\int_0^{\infty} \frac{x^{1-s}}{1+x^2} dx = \frac{\pi}{2 \sin\left(\frac{\pi s}{2}\right)},$$

we get

$$\frac{\pi}{2} \frac{1}{2\pi i} \int_{(c)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) z^{-s} ds = \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + z^2} du + \frac{\pi}{2} \int_0^{\infty} \frac{u \sin(u)}{u^2 + z^2} du. \quad (3.35)$$

From [26, p. 65, Formula (15)],

$$\int_0^{\infty} \frac{u \sin(u)}{u^2 + z^2} du = \frac{\pi}{2} e^{-z}. \quad (3.36)$$

Substituting (3.36) in (3.35) and multiplying both sides by $2/\pi$, we arrive at (3.32). The identity also holds, by analytic continuation, for $\text{Re}(z) > 0$ since both sides are analytic in this region.

Now again, shifting the line of integration from $\operatorname{Re}(s) = c$ to $\operatorname{Re}(s) = d$, $1 < d < 2$, noting that there is no pole of the integrand, applying Cauchy's residue theorem and making use of the fact that the integrals along the horizontal segments tend to zero as the height of the contour tends to ∞ , we see that for $\operatorname{Re}(w) > 0$,

$$\frac{1}{2\pi i} \int_{(c)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) (2\pi m w)^{-s} ds = \frac{1}{2\pi i} \int_{(d)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) (2\pi m w)^{-s} ds.$$

Hence, from (3.32) with $z = 2\pi m w$,

$$\sum_{m=1}^{\infty} \frac{1}{2\pi i} \int_{(d)} \Gamma(s) \psi(s) \cot\left(\frac{\pi s}{2}\right) (2\pi m w)^{-s} ds = \sum_{m=1}^{\infty} \left\{ \frac{2}{\pi} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + (2\pi m w)^2} du + \frac{\pi}{2} e^{-2\pi m w} \right\}. \quad (3.37)$$

Therefore, substituting (3.37) and (3.31) in (3.27), we get

$$\begin{aligned} J_1 &= 2\pi \left[\frac{(\gamma - \log(w))}{2\pi} (2 \log(w) - (\psi(iw) + \psi(-iw))) + \sum_{m=1}^{\infty} \left\{ \frac{2}{\pi} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + (2\pi m w)^2} du + \frac{\pi}{2} e^{-2\pi m w} \right\} \right] \\ &= 2(\gamma - \log(w)) \left(\log(w) - \frac{1}{2} (\psi(iw) + \psi(-iw)) \right) + 4 \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + (2\pi m w)^2} du + \frac{\pi^2}{e^{2\pi w} - 1}. \end{aligned} \quad (3.38)$$

Now from (3.24), (3.26) and (3.38), we have

$$\begin{aligned} &\psi_1(iw + 1) - \frac{1}{2} \log^2(iw) + \psi_1(-iw + 1) - \frac{1}{2} \log^2(-iw) \\ &= 2(\gamma - \log(w)) \left(\log(w) - \frac{1}{2} (\psi(iw) + \psi(-iw)) \right) + 4 \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + (2\pi m w)^2} du + \frac{\pi^2}{e^{2\pi w} - 1} \\ &\quad - \pi^2 \left(\frac{1}{e^{2\pi w} - 1} - \frac{1}{2\pi w} \right). \end{aligned}$$

Now using (3.19) twice and using the elementary fact $\log(iw) - \log(-iw) = \pi i$ for $\operatorname{Re}(w) > 0$, we are led to

$$\begin{aligned} 4 \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u) \log(u)}{u^2 + (2\pi m w)^2} du &= \psi_1(iw) - \frac{1}{2} \log^2(iw) + \psi_1(-iw) - \frac{1}{2} \log^2(-iw) + \frac{\pi}{2w} \\ &\quad + (\gamma - \log(w)) (\psi(iw) + \psi(-iw) - 2 \log(w)). \end{aligned} \quad (3.39)$$

Lastly, employing (2.2) again with w replaced by $2\pi w$, we see that

$$-4 \log(w) \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u)}{u^2 + (2\pi m w)^2} du = \log(w) (\psi(iw) + \psi(-iw) - 2 \log(w)). \quad (3.40)$$

Finally, adding the respective sides of (3.39) and (3.40), we are led to (3.20). This completes the proof. \square

4. A NEW REPRESENTATION FOR $\left. \frac{\partial^2}{\partial b^2} E_{2,b}(z) \right|_{b=1}$

The two-variable Mittag-Leffler function $E_{\alpha,\beta}(z)$, introduced by Wiman [61], is defined by

$$E_{\alpha,\beta}(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)} \quad (\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0). \quad (4.1)$$

There is an extensive literature on these Mittag-Leffler functions, see, for example, [30], and the references therein. Yet, closed-form expressions exist only for the first derivatives of $E_{\alpha,\beta}(z)$ with respect to the parameters α and β . The reader is referred to a recent paper of Apelblat [1] for a collection of such evaluations most of which involve the Shi(z) and Chi(z) functions defined in (4.3).

From (4.1), it is easy to see that [1, Equation (98)]

$$\frac{\partial^2}{\partial \beta^2} E_{\alpha, \beta}(t) = \sum_{k=0}^{\infty} \frac{\psi^2(\alpha k + \beta) - \psi'(\alpha k + \beta)}{\Gamma(\alpha k + \beta)} t^k. \quad (4.2)$$

However, there are no closed-form evaluations known for the above series. In what follows, we establish a new result which transforms $\left. \frac{\partial^2}{\partial \beta^2} E_{2, \beta}(t) \right|_{\beta=1}$ into a suitable integral which is absolutely essential in proving Theorem 1.1. Before we embark upon the proof though, we need a few lemmas concerning the *hyperbolic sine and cosine integrals* Shi(z) and Chi(z) defined by [48, p. 150, Equation (6.2.15), (6.2.16)]

$$\text{Shi}(z) := \int_0^z \frac{\sinh(t)}{t} dt, \quad \text{Chi}(z) := \gamma + \log(z) + \int_0^z \frac{\cosh(t) - 1}{t} dt. \quad (4.3)$$

The first lemma involving these functions was established in [23, Lemma 9.1]. Here, and throughout the sequel, we use the following notation for brevity.

$$(\sinh \text{Shi} - \cosh \text{Chi})(w) := \sinh(w) \text{Shi}(w) - \cosh(w) \text{Chi}(w).$$

Lemma 4.1. *Let $\text{Re}(w) > 0$. Then*

$$\int_0^{\infty} \frac{t \cos t dt}{t^2 + w^2} = (\sinh \text{Shi} - \cosh \text{Chi})(w).$$

We require another result from [22, Lemma 3.2].

Lemma 4.2. *For $\text{Re}(w) > 0$,*

$$\sum_{k=0}^{\infty} \frac{\psi(2k+1)}{\Gamma(2k+1)} w^{2k} = (\sinh \text{Shi} - \cosh \text{Chi})(w) + \log(w) \cosh(w).$$

We are now ready to prove our result on evaluating the second derivative of the Mittag-Leffler function $E_{2,b}(z)$ with respect to the parameter b at $b = 1$.

Theorem 4.3. *Let $b > 0$ and $\text{Re}(w) > 0$. Let the two-variable Mittag-Leffler function $E_{2,b}(z)$ be defined in (4.1). Then*

$$\left. \frac{\partial^2}{\partial b^2} E_{2,b}(w^2) \right|_{b=1} = \log^2(w) \cosh(w) + 2 \int_0^{\infty} \frac{u \cos(u) \log(u) du}{u^2 + w^2}. \quad (4.4)$$

Proof. To prove this result, we first apply a result of Dzhrbashyan [25, p. 130, Equation (2.12)] that will be used in the proof. It states that for $0 \leq \arg(z) < \pi$ or $-\pi < \arg(z) \leq 0$, one has⁷

$$E_{2,b}(z) = \frac{1}{2} z^{\frac{1}{2}(1-b)} \left\{ e^{\sqrt{z}} + e^{\mp i\pi(1-b) - \sqrt{z}} \right\} + \frac{\sin(\pi b)}{2\pi} \int_0^{\infty} \frac{e^{\pm i(\sqrt{t} - \frac{\pi}{2}(1-b))}}{t+z} t^{\frac{1}{2}(1-b)} dt, \quad (4.5)$$

where the upper or lower signs are taken, respectively, for $0 \leq \arg(z) < \pi$ or $-\pi < \arg(z) \leq 0$.

We first assume $0 \leq \arg(z) < \pi$. Differentiating (4.5) twice with respect to b and simplifying leads to

$$\begin{aligned} \frac{\partial^2}{\partial b^2} E_{2,b}(z) &= \frac{1}{2} \left[-\frac{1}{2} \log(z) \left\{ -\frac{1}{2} z^{\frac{1}{2}(1-b)} \log(z) \left(e^{\sqrt{z}} + e^{-i\pi(1-b) - \sqrt{z}} \right) + i\pi z^{\frac{1}{2}(1-b)} e^{i\pi(b-1) - \sqrt{z}} \right\} \right. \\ &\quad \left. + i\pi \sqrt{z} e^{-\sqrt{z}} z^{-\frac{b}{2}} e^{i\pi(b-1)} \left(i\pi - \frac{1}{2} \log(z) \right) \right] - \frac{\pi \sin(\pi b)}{2} \int_0^{\infty} \frac{e^{i(\sqrt{t} - \frac{\pi}{2}(1-b))}}{t+z} t^{\frac{1}{2}(1-b)} dt \\ &\quad + \frac{1}{2} \cos(\pi b) \int_0^{\infty} \frac{e^{i(\sqrt{t} - \frac{\pi}{2}(1-b))}}{t+z} t^{\frac{1}{2}(1-b)} (i\pi - \log(t)) dt \end{aligned}$$

⁷It is to be noted that Dzhrbashyan uses a different notation for the two-variable Mittag-Leffler function in his book, namely, $E_{\rho}(z; b) = \sum_{k=0}^{\infty} z^k / \Gamma(b + k\rho^{-1})$. See [25, p. 117]. We have used the contemporary notation here.

$$+ \frac{1}{8\pi} \sin(\pi b) \int_0^\infty \frac{e^{i(\sqrt{t}-\frac{\pi}{2}(1-b))}}{t+z} t^{\frac{1}{2}(1-b)} (i\pi - \log(t))^2 dt,$$

so that

$$\frac{\partial^2}{\partial b^2} E_{2,b}(z) \Big|_{b=1} = \frac{1}{4} \log^2(z) \cosh(\sqrt{z}) - \frac{i\pi}{2} e^{-\sqrt{z}} \log(z) - \frac{\pi^2}{2} e^{-\sqrt{z}} - \frac{1}{2} \int_0^\infty \frac{e^{i\sqrt{t}}(i\pi - \log(t))}{t+z} dt. \quad (4.6)$$

Now employing the change of variable $t = u^2$, we have

$$\begin{aligned} \int_0^\infty \frac{e^{i\sqrt{t}}(i\pi - \log(t))}{t+z} dt &= 2 \int_0^\infty \frac{ue^{iu}(i\pi - 2\log(u))}{u^2+z} du \\ &= 2\pi i \int_0^\infty \frac{u \cos(u)}{u^2+z} du - 2\pi \int_0^\infty \frac{u \sin(u)}{u^2+z} du - 4 \int_0^\infty \frac{ue^{iu} \log(u)}{u^2+z} du. \end{aligned} \quad (4.7)$$

Next, from Lemma 4.1,

$$\int_0^\infty \frac{u \cos(u)}{u^2+z} du = (\sinh \operatorname{Shi} - \cosh \operatorname{Chi})(\sqrt{z}). \quad (4.8)$$

Moreover,

$$\int_0^\infty \frac{ue^{iu} \log(u)}{u^2+z} du = \int_0^\infty \frac{u \cos(u) \log(u)}{u^2+z} du + i \int_0^\infty \frac{u \sin(u) \log(u)}{u^2+z} du, \quad (4.9)$$

Now from [50, p. 537, Formula **2.6.32.8**]⁸,

$$\int_0^\infty \frac{u \sin(u) \log(u)}{u^2+z} du = \frac{\pi}{2} \left\{ e^{-\sqrt{z}} \log(\sqrt{z}) - \frac{1}{2} \left(e^{\sqrt{z}} \operatorname{Ei}(-\sqrt{z}) + e^{-\sqrt{z}} \operatorname{Ei}(\sqrt{z}) \right) \right\}. \quad (4.10)$$

Hence along with Lemma 4.1, (3.30) and (4.10) imply

$$\int_0^\infty \frac{u \sin(u) \log(u)}{u^2+z} du = \frac{\pi}{2} \left\{ e^{-\sqrt{z}} \log(\sqrt{z}) + (\sinh \operatorname{Shi} - \cosh \operatorname{Chi})(\sqrt{z}) \right\}. \quad (4.11)$$

Therefore, from (4.7), (4.8), (3.36), (4.9) and (4.11), we arrive at

$$\int_0^\infty \frac{e^{i\sqrt{t}}(i\pi - \log(t))}{t+z} dt = -\pi^2 e^{-\sqrt{z}} - 2\pi i e^{-\sqrt{z}} \log(\sqrt{z}) - 4 \int_0^\infty \frac{u \cos(u) \log(u)}{u^2+z} du. \quad (4.12)$$

Finally, substituting (4.12) in (4.6), we arrive at

$$\frac{\partial^2}{\partial b^2} E_{2,b}(z) \Big|_{b=1} = \log^2(\sqrt{z}) \cosh(\sqrt{z}) + 2 \int_0^\infty \frac{u \cos(u) \log(u)}{u^2+z} du.$$

Now let $z = w^2$ to arrive at (4.4) for $0 \leq \arg(w) < \frac{\pi}{2}$. Similarly starting with $-\pi < \arg(z) \leq 0$ and proceeding exactly as above and putting $z = w^2$ at the end, we again obtain (4.4) for $-\frac{\pi}{2} < \arg(w) \leq 0$. Hence we see that (4.4) actually holds for $\operatorname{Re}(w) > 0$. \square

5. PROOF OF THE MAIN RESULTS

This section is devoted to proving the transformation for the Lambert series for logarithm given in Theorem 1.1, the asymptotic expansion of this series in Theorem 1.2, and a mean value theorem in Theorem 1.3.

⁸The formula given in the book has a typo. We have corrected it here.

5.1. Proof of Theorem 1.1. The idea is to differentiate both sides of (1.5) with respect to a and then let $a \rightarrow 0$. Define

$$\begin{aligned} F_1(a, y) &:= \frac{d}{da} \sum_{n=1}^{\infty} \sigma_a(n) e^{-ny}, \\ F_2(a, y) &:= \frac{d}{da} \left\{ \frac{1}{2} \left(\left(\frac{2\pi}{y} \right)^{1+a} \operatorname{cosec} \left(\frac{\pi a}{2} \right) + 1 \right) \zeta(-a) - \frac{1}{y} \zeta(1-a) \right\}, \\ F_3(a, y) &:= \frac{d}{da} \left\{ \frac{2\pi}{y \sin \left(\frac{\pi a}{2} \right)} \sum_{n=1}^{\infty} \sigma_a(n) \left(\frac{(2\pi n)^{-a}}{\Gamma(1-a)} {}_1F_2 \left(1; \frac{1-a}{2}, 1 - \frac{a}{2}; \frac{4\pi^4 n^2}{y^2} \right) - \left(\frac{2\pi}{y} \right)^a \cosh \left(\frac{4\pi^2 n}{y} \right) \right) \right\}, \end{aligned} \quad (5.1)$$

and let

$$G_1(y) = \lim_{a \rightarrow 0} F_1(a, y), \quad G_2(y) = \lim_{a \rightarrow 0} F_2(a, y), \quad G_3(y) = \lim_{a \rightarrow 0} F_3(a, y). \quad (5.2)$$

Clearly, from (1.5),

$$G_1(y) + G_2(y) = G_3(y). \quad (5.3)$$

Using (1.3), it is readily seen that

$$G_1(y) = \sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1}. \quad (5.4)$$

By routine differentiation,

$$\begin{aligned} F_2(a, y) &= \frac{1}{y} \zeta'(1-a) - \frac{1}{2} \zeta'(-a) - \frac{1}{2} \left(\frac{2\pi}{y} \right)^{1+a} \operatorname{cosec} \left(\frac{\pi a}{2} \right) \zeta'(-a) \\ &\quad + \frac{1}{2} \zeta(-a) \left(\frac{2\pi}{y} \right)^{1+a} \operatorname{cosec} \left(\frac{\pi a}{2} \right) \left(\log \left(\frac{2\pi}{y} \right) - \frac{\pi}{2} \cot \left(\frac{\pi a}{2} \right) \right). \end{aligned} \quad (5.5)$$

Letting $a \rightarrow 0$ in (5.5) leads to

$$G_2(y) = \frac{1}{4} \log(2\pi) - \frac{1}{2y} \log^2(y) + \frac{\gamma^2}{2y} - \frac{\pi^2}{12y}, \quad (5.6)$$

which is now proved. Note that

$$\begin{aligned} G_2(y) &= \lim_{a \rightarrow 0} \left(-\frac{1}{2} \zeta'(-a) \right) \\ &\quad + \lim_{a \rightarrow 0} \left[\frac{1}{y} \zeta'(1-a) - \frac{(2\pi/y)^{1+a}}{2 \sin \left(\frac{\pi a}{2} \right)} \left\{ \zeta'(-a) - \zeta(-a) \log \left(\frac{2\pi}{y} \right) + \frac{\pi}{2} \zeta(-a) \cot \left(\frac{\pi a}{2} \right) \right\} \right]. \end{aligned} \quad (5.7)$$

We make use of the following well-known Laurent series expansions around $a = 0$:

$$\begin{aligned} \zeta(-a) &= -\frac{1}{2} + \frac{1}{2} \log(2\pi) a + \left[\frac{\gamma^2}{4} - \frac{\pi^2}{48} - \frac{\log^2(2\pi)}{4} + \frac{\gamma_1}{2} \right] a^2 + O(a^3), \\ \zeta'(1-a) &= -\frac{1}{a^2} - \gamma_1 - \gamma_2 a + O(a^2), \\ \left(\frac{2\pi}{y} \right)^{1+a} &= \frac{2\pi}{y} + \frac{2\pi}{y} \log \left(\frac{2\pi}{y} \right) a + \frac{\pi}{y} \log^2 \left(\frac{2\pi}{y} \right) a^2 + O(a^3), \\ \operatorname{cosec} \left(\frac{\pi a}{2} \right) &= \frac{2}{\pi a} + \frac{\pi}{12} a + O(a^3), \\ \cot \left(\frac{\pi a}{2} \right) &= \frac{2}{\pi a} - \frac{\pi}{6} a + O(a^3). \end{aligned} \quad (5.8)$$

Hence substituting (5.8) in (5.7), we get

$$\begin{aligned}
G_2(y) &= \frac{1}{4} \log(2\pi) + \lim_{a \rightarrow 0} \left[-\frac{1}{ya^2} - \frac{\gamma_1}{y} + O(a) + \left\{ -\frac{2}{ya} - \frac{2}{y} \log\left(\frac{2\pi}{y}\right) + \left(-\frac{\pi^2}{12y} - \frac{1}{y} \log^2\left(\frac{2\pi}{y}\right)\right) a + O(a^2) \right\} \right. \\
&\quad \left. \times \left\{ -\frac{1}{2a} + \frac{1}{2} \log\left(\frac{2\pi}{y}\right) + \left(-\frac{\gamma^2}{4} + \frac{\pi^2}{16} - \frac{\gamma_1}{2} + \frac{1}{4} \log^2(2\pi) - \frac{1}{2} \log(2\pi) \log\left(\frac{2\pi}{y}\right)\right) a + O(a^2) \right\} \right] \\
&= \frac{1}{4} \log(2\pi) + \lim_{a \rightarrow 0} \left[\left(-\frac{1}{y} + \frac{1}{y}\right) \frac{1}{a^2} + \left(-\frac{1}{y} \log\left(\frac{2\pi}{y}\right) + \frac{1}{y} \log\left(\frac{2\pi}{y}\right)\right) \frac{1}{a} \right. \\
&\quad \left. + \left(-\frac{\gamma_1}{y} + \frac{\gamma^2}{2y} - \frac{\pi^2}{12y} + \frac{\gamma_1}{y} - \frac{1}{2y} \left\{ \log^2(2\pi) - 2 \log(2\pi) \log\left(\frac{2\pi}{y}\right) + \log^2\left(\frac{2\pi}{y}\right) \right\}\right) + O(a) \right] \\
&= \frac{1}{4} \log(2\pi) + \frac{\gamma^2}{2y} - \frac{\pi^2}{12y} - \frac{\log^2 y}{2y}.
\end{aligned}$$

Next, we apply the definitions of $\sigma_a(n)$, ${}_1F_2$ and simplify to obtain

$$\begin{aligned}
F_3(a, y) &= \frac{d}{da} \left[\frac{2\pi}{y \sin\left(\frac{\pi a}{2}\right)} \sum_{m, n=1}^{\infty} \left\{ (2\pi n)^{-a} \sum_{k=0}^{\infty} \frac{(4\pi^2 mn/y)^{2k}}{\Gamma(1-a+2k)} - \left(\frac{2\pi m}{y}\right)^a \cosh\left(\frac{4\pi^2 mn}{y}\right) \right\} \right] \\
&= \frac{2\pi}{y} \sum_{m, n=1}^{\infty} \sum_{k=0}^{\infty} \left(\frac{4\pi^2 mn}{y}\right)^{2k} \frac{1}{\sin^2\left(\frac{\pi a}{2}\right)} \left[\frac{(2\pi n)^{-a}}{\Gamma(1-a+2k)} \left\{ (\psi(1-a+2k) - \log(2\pi n)) \sin\left(\frac{\pi a}{2}\right) \right. \right. \\
&\quad \left. \left. - \frac{\pi}{2} \cos\left(\frac{\pi a}{2}\right) \right\} - \frac{1}{\Gamma(2k+1)} \left(\frac{2\pi m}{y}\right)^a \left(\log\left(\frac{2\pi m}{y}\right) \sin\left(\frac{\pi a}{2}\right) - \frac{\pi}{2} \cos\left(\frac{\pi a}{2}\right) \right) \right] \\
&= \frac{2\pi}{y} \sum_{m, n=1}^{\infty} \sum_{k=0}^{\infty} \left(\frac{4\pi^2 mn}{y}\right)^{2k} \frac{A_1(a) - A_2(a)}{\sin^2\left(\frac{\pi a}{2}\right)},
\end{aligned}$$

where in the first step we have interchanged the order of summation and differentiation using the fact [23, p. 35] that the series in the definition of (5.1) converges uniformly as long as $\text{Re}(a) > -1$, and where

$$\begin{aligned}
A_1(a) &= A_1(m, n, k; a) := \frac{(2\pi n)^{-a}}{\Gamma(1-a+2k)} \left\{ (\psi(1-a+2k) - \log(2\pi n)) \sin\left(\frac{\pi a}{2}\right) - \frac{\pi}{2} \cos\left(\frac{\pi a}{2}\right) \right\}, \\
A_2(a) &= A_2(m, n, k; a) := \frac{1}{\Gamma(2k+1)} \left(\frac{2\pi m}{y}\right)^a \left\{ \log\left(\frac{2\pi m}{y}\right) \sin\left(\frac{\pi a}{2}\right) - \frac{\pi}{2} \cos\left(\frac{\pi a}{2}\right) \right\}.
\end{aligned}$$

We now show that

$$\begin{aligned}
\lim_{a \rightarrow 0} \frac{A_1(a) - A_2(a)}{\sin^2\left(\frac{\pi a}{2}\right)} &= \frac{1}{\pi \Gamma(2k+1)} \left\{ \log\left(\frac{4\pi^2 mn}{y}\right) \log\left(\frac{ny}{m}\right) - 2 \log(2n\pi) \psi(2k+1) + \psi^2(2k+1) \right. \\
&\quad \left. - \psi'(2k+1) \right\}. \tag{5.9}
\end{aligned}$$

Let L denote the limit in the equation given above. The expression inside the limit is of the form $0/0$. By routine calculation, we find that

$$A'_1(a) = \frac{(2\pi n)^{-a}}{\Gamma(1-a+2k)} \left\{ -\psi'(1-a+2k) + \frac{\pi^2}{4} + (\psi(1-a+2k) - \log(2\pi n))^2 \right\} \sin\left(\frac{\pi a}{2}\right), \tag{5.10}$$

$$\begin{aligned}
A'_2(a) &= \frac{(2\pi m/y)^a}{\Gamma(2k+1)} \left\{ \log\left(\frac{2\pi m}{y}\right) \left(\log\left(\frac{2\pi m}{y}\right) \sin\left(\frac{\pi a}{2}\right) - \frac{\pi}{2} \cos\left(\frac{\pi a}{2}\right) \right) \right. \\
&\quad \left. + \frac{\pi}{2} \log\left(\frac{2\pi m}{y}\right) \cos\left(\frac{\pi a}{2}\right) + \frac{\pi^2}{4} \sin\left(\frac{\pi a}{2}\right) \right\}, \tag{5.11}
\end{aligned}$$

where ' denotes differentiation with respect to a . Since $A_1'(0) = A_2'(0) = 0$, using L'Hopital's rule again, it is seen that

$$L = \frac{A_1''(0) - A_2''(0)}{\pi^2/2},$$

Differentiating (5.10) and (5.11) with respect to a , we get

$$\begin{aligned} A_1''(0) &= \frac{\pi}{2\Gamma(2k+1)} \left\{ -\psi'(2k+1) + \frac{\pi^2}{4} + (\psi(2k+1) - \log(2\pi n))^2 \right\}, \\ A_2''(0) &= \frac{1}{\Gamma(2k+1)} \left\{ \frac{\pi}{2} \log^2 \left(\frac{2\pi m}{y} \right) + \frac{\pi^3}{8} \right\}, \end{aligned}$$

whence we obtain (5.9) upon simplification. Therefore, from (5.2) and (5.9), we deduce that

$$\begin{aligned} G_3(y) &= \frac{2}{y} \sum_{m,n=1}^{\infty} \sum_{k=0}^{\infty} \frac{(4\pi^2 mn/y)^{2k}}{(2k)!} \left\{ \log \left(\frac{4\pi^2 mn}{y} \right) \log \left(\frac{ny}{m} \right) - 2 \log(2n\pi) \psi(2k+1) \right. \\ &\quad \left. + \psi^2(2k+1) - \psi'(2k+1) \right\} \\ &= \frac{2}{y} \sum_{m,n=1}^{\infty} \left\{ \log \left(\frac{4\pi^2 mn}{y} \right) \log \left(\frac{ny}{m} \right) \cosh \left(\frac{4\pi^2 mn}{y} \right) - 2 \log(2n\pi) \sum_{k=0}^{\infty} \frac{\psi(2k+1)}{\Gamma(2k+1)} \left(\frac{4\pi^2 mn}{y} \right)^{2k} \right. \\ &\quad \left. + \sum_{k=0}^{\infty} \frac{\psi^2(2k+1) - \psi'(2k+1)}{\Gamma(2k+1)} \left(\frac{4\pi^2 mn}{y} \right)^{2k} \right\}. \end{aligned}$$

Therefore, invoking Lemma 4.2, using (4.2) and Theorem 4.3 with $w = 4\pi^2 mn/y$, we are led to

$$\begin{aligned} G_3(y) &= \frac{2}{y} \sum_{m,n=1}^{\infty} \left\{ \log \left(\frac{4\pi^2 mn}{y} \right) \log \left(\frac{ny}{m} \right) \cosh \left(\frac{4\pi^2 mn}{y} \right) - 2 \log(2n\pi) \left[(\sinh \text{Shi} - \cosh \text{Chi}) \left(\frac{4\pi^2 mn}{y} \right) \right. \right. \\ &\quad \left. \left. + \log \left(\frac{4\pi^2 mn}{y} \right) \cosh \left(\frac{4\pi^2 mn}{y} \right) \right] + \log^2 \left(\frac{4\pi^2 mn}{y} \right) \cosh \left(\frac{4\pi^2 mn}{y} \right) + 2 \int_0^{\infty} \frac{u \cos(u) \log(u) du}{u^2 + (4\pi^2 mn/y)^2} \right\} \\ &= \frac{4}{y} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u) \log(u/(2\pi n))}{u^2 + (4\pi^2 mn/y)^2} du, \end{aligned}$$

where in the last step, we used Lemma 4.1 with $w = 4\pi^2 mn/y$.

Next, an application of Theorem 3.4 with $w = 2\pi n/y$ and then of (2.2) with $w = 4\pi^2 n/y$ in the second step further yields

$$\begin{aligned} G_3(y) &= \frac{4}{y} \sum_{n=1}^{\infty} \left\{ \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u) \log(uy/(2\pi n))}{u^2 + (4\pi^2 mn/y)^2} du - \log(y) \sum_{m=1}^{\infty} \int_0^{\infty} \frac{u \cos(u)}{u^2 + (4\pi^2 mn/y)^2} du \right\} \\ &= \frac{1}{y} \sum_{n=1}^{\infty} \left\{ \psi_1 \left(\frac{2\pi in}{y} \right) - \frac{1}{2} \log^2 \left(\frac{2\pi in}{y} \right) + \psi_1 \left(-\frac{2\pi in}{y} \right) - \frac{1}{2} \log^2 \left(-\frac{2\pi in}{y} \right) + \frac{y}{4n} \right. \\ &\quad \left. + (\gamma + \log(y)) \left(\psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) - 2 \log \left(\frac{2\pi n}{y} \right) \right) \right\} \\ &= \frac{1}{y} \sum_{n=1}^{\infty} \left\{ \psi_1 \left(\frac{2\pi in}{y} \right) - \frac{1}{2} \log^2 \left(\frac{2\pi in}{y} \right) + \psi_1 \left(-\frac{2\pi in}{y} \right) - \frac{1}{2} \log^2 \left(-\frac{2\pi in}{y} \right) + \frac{y}{4n} \right\} \\ &\quad + \frac{(\gamma + \log(y))}{y} \sum_{n=1}^{\infty} \left\{ \left(\psi \left(\frac{2\pi in}{y} \right) + \psi \left(-\frac{2\pi in}{y} \right) - 2 \log \left(\frac{2\pi n}{y} \right) \right) \right\}, \end{aligned} \tag{5.12}$$

where the validity of the last step results from the fact that invoking Theorem 3.2, once with $z = 2\pi in/y$, and again with $z = -2\pi in/y$, and adding the two expansions yields

$$\psi_1\left(\frac{2\pi in}{y}\right) - \frac{1}{2}\log^2\left(\frac{2\pi in}{y}\right) + \psi_1\left(-\frac{2\pi in}{y}\right) - \frac{1}{2}\log^2\left(-\frac{2\pi in}{y}\right) + \frac{y}{4n} = O_y\left(\frac{\log(n)}{n^2}\right). \quad (5.13)$$

Therefore, from (5.3), (5.4), (5.6) and (5.12), we arrive at

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1} + \frac{1}{4}\log(2\pi) - \frac{1}{2y}\log^2(y) + \frac{\gamma^2}{2y} - \frac{\pi^2}{12y} \\ &= \frac{1}{y} \sum_{n=1}^{\infty} \left\{ \psi_1\left(\frac{2\pi in}{y}\right) - \frac{1}{2}\log^2\left(\frac{2\pi in}{y}\right) + \psi_1\left(-\frac{2\pi in}{y}\right) - \frac{1}{2}\log^2\left(-\frac{2\pi in}{y}\right) + \frac{y}{4n} \right\} \\ & \quad + \frac{(\gamma + \log(y))}{y} \sum_{n=1}^{\infty} \left\{ \left(\psi\left(\frac{2\pi in}{y}\right) + \psi\left(-\frac{2\pi in}{y}\right) - 2\log\left(\frac{2\pi n}{y}\right) \right) \right\}, \end{aligned}$$

which, upon rearrangement, gives (1.13).

Now (1.14) can be immediately derived from (1.13) by invoking (1.6) to transform the first series on the right-hand side of (1.13), multiplying both sides of the resulting identity by y followed by simplification and rearrangement. \square

The theorem just proved allows us to obtain the complete asymptotic expansion of $\sum_{n=1}^{\infty} d(n)\log(n)e^{-ny}$ as $y \rightarrow 0$. This is derived next.

5.2. Proof of Theorem 1.2. From (3.13), as $y \rightarrow 0$,

$$\psi\left(\frac{2\pi in}{y}\right) \sim \log\left(\frac{2\pi in}{y}\right) - \sum_{k=1}^{\infty} \frac{B_{2k}}{2k} \left(\frac{2\pi in}{y}\right)^{-2k} \quad \text{and} \quad \psi\left(-\frac{2\pi in}{y}\right) \sim \log\left(-\frac{2\pi in}{y}\right) - \sum_{k=1}^{\infty} \frac{B_{2k}}{2k} \left(-\frac{2\pi in}{y}\right)^{-2k}.$$

Inserting the above asymptotics into the first sum of the right-hand side of (1.13), we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \left\{ \log\left(\frac{2\pi n}{y}\right) - \frac{1}{2} \left(\psi\left(\frac{2\pi in}{y}\right) + \psi\left(-\frac{2\pi in}{y}\right) \right) \right\} &\sim \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{(-1)^k B_{2k}}{2k} \left(\frac{2\pi n}{y}\right)^{-2k} \\ &= \sum_{k=1}^{\infty} \frac{(-1)^k B_{2k} \zeta(2k)}{2k} \left(\frac{2\pi}{y}\right)^{-2k} \\ &= -\frac{1}{2} \sum_{k=1}^{\infty} \frac{B_{2k}^2 y^{2k}}{(2k)(2k)!}, \end{aligned} \quad (5.14)$$

where in the last step we used Euler's formula

$$\zeta(2m) = (-1)^{m+1} \frac{(2\pi)^{2m} B_{2m}}{2(2m)!} \quad (m \geq 0). \quad (5.15)$$

Next, we need to find the asymptotic expansion of the second sum on the right-hand side of (1.13). Invoking Theorem 3.2 twice, once with $z = 2\pi in/y$, and again, with $z = -2\pi in/y$, and inserting them into the second sum of the right-hand side of (1.13), we obtain

$$\begin{aligned} & \sum_{n=1}^{\infty} \left\{ \psi_1\left(\frac{2\pi in}{y}\right) + \psi_1\left(-\frac{2\pi in}{y}\right) - \frac{1}{2} \left(\log^2\left(\frac{2\pi in}{y}\right) + \log^2\left(-\frac{2\pi in}{y}\right) \right) + \frac{y}{4n} \right\} \\ & \sim \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{(-1)^k B_{2k}}{k} \left(\frac{y}{2\pi n}\right)^{2k} \left[\frac{1}{2k-1} + \sum_{j=1}^{2k-2} \frac{1}{j} - \log\left(\frac{2\pi n}{y}\right) \right] \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{(-1)^k B_{2k}}{k} \left(\frac{y}{2\pi n}\right)^{2k} \left[\frac{1}{2k-1} + \left(\sum_{j=1}^{2k-2} \frac{1}{j} - \log\left(\frac{2\pi}{y}\right) \right) - \log n \right] \\
&= \sum_{k=1}^{\infty} \frac{(-1)^k B_{2k}}{k} \left(\frac{y}{2\pi}\right)^{2k} \left[\zeta(2k) \left\{ \frac{1}{2k-1} + \left(\sum_{j=1}^{2k-2} \frac{1}{j} - \log\left(\frac{2\pi}{y}\right) \right) \right\} + \zeta'(2k) \right] \\
&= - \sum_{k=1}^{\infty} \left[\frac{B_{2k}^2 y^{2k}}{(2k-1)(2k)(2k)!} + \frac{B_{2k}^2 y^{2k}}{2k(2k)!} \left(\sum_{j=1}^{2k-2} \frac{1}{j} - \log\left(\frac{2\pi}{y}\right) \right) + \frac{(-1)^{k+1} B_{2k} \zeta'(2k)}{k} \left(\frac{y}{2\pi}\right)^{2k} \right], \quad (5.16)
\end{aligned}$$

where in the penultimate step we used $\zeta'(s) = -\sum_{n=1}^{\infty} \log(n)n^{-s}$ for $\text{Re}(s) > 1$, and in the last step we used (5.15). Substituting (5.14) and (5.16) in (1.13), we see that as $y \rightarrow 0$,

$$\begin{aligned}
&\sum_{n=1}^{\infty} \frac{\log(n)}{e^{ny} - 1} \\
&\sim -\frac{1}{4} \log(2\pi) + \frac{1}{2y} \log^2(y) - \frac{\gamma^2}{2y} + \frac{\pi^2}{12y} - \frac{2}{y}(\gamma + \log(y)) \left\{ -\frac{1}{2} \sum_{k=1}^{\infty} \frac{B_{2k}^2 y^{2k}}{(2k)(2k)!} \right\} \\
&\quad - \frac{1}{y} \sum_{k=1}^{\infty} \left[\frac{B_{2k}^2 y^{2k}}{(2k-1)(2k)(2k)!} + \frac{B_{2k}^2 y^{2k}}{2k(2k)!} \left(\sum_{j=1}^{2k-2} \frac{1}{j} - \log\left(\frac{2\pi}{y}\right) \right) + \frac{(-1)^{k+1} B_{2k} \zeta'(2k)}{k} \left(\frac{y}{2\pi}\right)^{2k} \right] \\
&= -\frac{1}{4} \log(2\pi) + \frac{1}{2y} \log^2(y) - \frac{\gamma^2}{2y} + \frac{\pi^2}{12y} + \sum_{k=1}^{\infty} \frac{B_{2k} y^{2k-1}}{k} \left\{ \frac{B_{2k}}{2(2k)!} \left(\gamma - \sum_{j=1}^{2k-1} \frac{1}{j} + \log(2\pi) \right) + \frac{(-1)^k \zeta'(2k)}{(2\pi)^{2k}} \right\} \\
&= -\frac{1}{4} \log(2\pi) + \frac{1}{2y} \log^2(y) - \frac{\gamma^2}{2y} + \frac{\pi^2}{12y} + \frac{y}{6} \left\{ \frac{1}{24}(\gamma - 1 + \log(2\pi)) - \frac{1}{4\pi^2} \left[\frac{\pi^2}{6}(\gamma + \log(2\pi)) \right. \right. \\
&\quad \left. \left. - 12 \log(A) \right] \right\} + \sum_{k=2}^{\infty} \frac{B_{2k} y^{2k-1}}{k} \left\{ \frac{B_{2k}}{2(2k)!} \left(\gamma - \sum_{j=1}^{2k-1} \frac{1}{j} + \log(2\pi) \right) + \frac{(-1)^k \zeta'(2k)}{(2\pi)^{2k}} \right\} \\
&= \frac{1}{2y} \log^2(y) + \frac{1}{y} \left(\frac{\pi^2}{12} - \frac{\gamma^2}{2} \right) - \frac{1}{4} \log(2\pi) + \frac{y}{12} \left(\log(A) - \frac{1}{12} \right) \\
&\quad + \sum_{k=2}^{\infty} \frac{B_{2k} y^{2k-1}}{k} \left\{ \frac{B_{2k}}{2(2k)!} \left(\gamma - \sum_{j=1}^{2k-1} \frac{1}{j} + \log(2\pi) \right) + \frac{(-1)^k \zeta'(2k)}{(2\pi)^{2k}} \right\}.
\end{aligned}$$

This completes the proof. \square

An application of the result proved above in the theory of the moments of the Riemann zeta function is given next.

5.3. Proof of Theorem 1.3. Our goal is to link $\int_0^{\infty} \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt$ with $\sum_{n=1}^{\infty} \frac{\log(n)}{\exp(2\pi i n e^{-i\delta}) - 1}$ and then invoke Theorem 1.2. Our treatment is similar to that of Atkinson [3].

First note that, for $\text{Re}(y) > 0$,

$$\sum_{n=1}^{\infty} d(n) \log(n) e^{-ny} = \sum_{n=1}^{\infty} d(n) \log(n) \frac{1}{2\pi i} \int_{(c)} \Gamma(s) (ny)^{-s} ds, \quad (5.17)$$

where $c = \operatorname{Re}(s) > 1$. Using $(1 * \log)(n) = \sum_{d|n} \log d = \frac{1}{2}d(n) \log(n)$ and interchanging the order of summation and integration on the right-hand side of (5.17), we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} d(n) \log(n) e^{-ny} &= \frac{1}{2\pi i} \int_{(c)} \Gamma(s) \left(\sum_{n=1}^{\infty} \frac{d(n) \log(n)}{n^s} \right) y^{-s} ds \\ &= \frac{2}{2\pi i} \int_{(c)} \Gamma(s) \sum_{n=1}^{\infty} \frac{(1 * \log)(n)}{n^s} y^{-s} ds \\ &= -\frac{2}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) \zeta'(s) y^{-s} ds. \end{aligned} \quad (5.18)$$

Letting $y = 2\pi i e^{-i\delta}$ so that $\operatorname{Re}(\delta) > 0$, we deduce that

$$\sum_{n=1}^{\infty} d(n) \log(n) \exp(-2\pi i n e^{-i\delta}) = -\frac{2}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) \zeta'(s) (2\pi i e^{-i\delta})^{-s} ds. \quad (5.19)$$

We would like to shift the line of integration from $\operatorname{Re}(s) = c$ to $\operatorname{Re}(s) = 1/2$. The integrand in (5.19) has a third order pole at $s = 1$. Therefore, using Cauchy's residue theorem on the rectangular contour with sides $[c - iT, c + iT]$, $[c + iT, \frac{1}{2} + iT]$, $[\frac{1}{2} + iT, \frac{1}{2} - iT]$ and $[\frac{1}{2} - iT, c - iT]$, where $T > 0$, noting that (2.1) implies that the integrals along the horizontal line segments tend to zero as $T \rightarrow \infty$, we obtain

$$\begin{aligned} &\frac{1}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) \zeta'(s) (2\pi i e^{-i\delta})^{-s} ds \\ &= \frac{1}{2\pi i} \int_{(\frac{1}{2})} \Gamma(s) \zeta(s) \zeta'(s) (2\pi i e^{-i\delta})^{-s} ds + \operatorname{Res}_{s=1} \Gamma(s) \zeta(s) \zeta'(s) (2\pi i e^{-i\delta})^{-s} \\ &= \frac{1}{4\pi i} \int_{(\frac{1}{2})} \frac{\zeta(1-s) \zeta'(s) (i e^{-i\delta})^{-s}}{\cos(\frac{\pi s}{2})} ds + \frac{1}{2\pi i e^{-i\delta}} \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12} - \frac{1}{2} \log^2(2\pi i e^{-i\delta}) \right), \end{aligned} \quad (5.20)$$

where, in the last step, we used the functional equation (3.25) with s replaced by $1-s$. Hence, from (5.19) and (5.20), we arrive at

$$\begin{aligned} \frac{e^{-\frac{i\delta}{2}}}{2i} \int_{(\frac{1}{2})} \frac{\zeta(1-s) \zeta'(s) e^{-is(\frac{\pi}{2}-\delta)}}{\cos(\frac{\pi s}{2})} ds &= -\pi e^{-\frac{i\delta}{2}} \sum_{n=1}^{\infty} d(n) \log(n) \exp(-2\pi i n e^{-i\delta}) \\ &\quad + \frac{i e^{\frac{i\delta}{2}}}{2} \left(\gamma^2 - \frac{\pi^2}{6} - \log^2(2\pi i e^{-i\delta}) \right). \end{aligned} \quad (5.21)$$

We next consider the difference of the integrals on the left-hand sides of (1.16) and (5.21) and employ the change of variable $s = \frac{1}{2} + it$ in the second integral in the first step below to see that

$$\begin{aligned} &\int_0^{\infty} \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt - \frac{e^{-\frac{i\delta}{2}}}{i} \int_{(\frac{1}{2})} \frac{\zeta(1-s) \zeta'(s)}{2 \cos(\frac{\pi s}{2})} e^{-is(\frac{\pi}{2}-\delta)} ds \\ &= \int_0^{\infty} \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt - e^{-\frac{i\delta}{2}} \int_{-\infty}^{\infty} \frac{\zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right)}{\left(e^{\frac{i\pi}{2}\left(\frac{1}{2}+it\right)} + e^{-\frac{i\pi}{2}\left(\frac{1}{2}+it\right)} \right)} e^{-i\left(\frac{1}{2}+it\right)\left(\frac{\pi}{2}-\delta\right)} dt \end{aligned}$$

$$\begin{aligned}
&= \int_0^\infty \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt + \int_0^\infty \frac{\zeta\left(\frac{1}{2} + it\right) \zeta'\left(\frac{1}{2} - it\right)}{\left(e^{\frac{i\pi}{4} + \frac{\pi t}{2}} + e^{-\frac{i\pi}{4} - \frac{\pi t}{2}}\right)} e^{-\frac{i\pi}{4} - \frac{\pi t}{2} + \delta t} dt \\
&\quad - \int_0^\infty \frac{\zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right)}{\left(e^{\frac{i\pi}{4} - \frac{\pi t}{2}} + e^{-\frac{i\pi}{4} + \frac{\pi t}{2}}\right)} e^{-\frac{i\pi}{4} + \frac{\pi t}{2} - \delta t} dt \\
&= \int_0^\infty \frac{\zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right)}{\left(e^{\frac{i\pi}{4} - \frac{\pi t}{2}} + e^{-\frac{i\pi}{4} + \frac{\pi t}{2}}\right)} e^{\frac{i\pi}{4} - \frac{\pi t}{2} - \delta t} dt + \int_0^\infty \frac{\zeta\left(\frac{1}{2} + it\right) \zeta'\left(\frac{1}{2} - it\right)}{\left(e^{\frac{i\pi}{4} + \frac{\pi t}{2}} + e^{-\frac{i\pi}{4} - \frac{\pi t}{2}}\right)} e^{-\frac{i\pi}{4} - \frac{\pi t}{2} + \delta t} dt. \tag{5.22}
\end{aligned}$$

From [29, p. 127, Eq. 20], we have

$$\zeta\left(\frac{1}{2} - it\right) \ll |t|^{\frac{1}{4} + \frac{\epsilon}{2}} \quad \text{and} \quad \zeta'\left(\frac{1}{2} + it\right) \ll |t|^{\frac{1}{4} + \frac{\epsilon}{2}},$$

whence, by reverse triangle inequality, we have

$$\left| \frac{\zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right)}{\left(e^{\frac{i\pi}{4} - \frac{\pi t}{2}} + e^{-\frac{i\pi}{4} + \frac{\pi t}{2}}\right)} e^{\frac{i\pi}{4} - \frac{\pi t}{2} - \delta t} \right| \ll \frac{|t|^{\frac{1}{2} + \epsilon} e^{-\operatorname{Re}(\delta)t - \frac{\pi t}{2}}}{e^{\frac{\pi t}{2}} - e^{-\frac{\pi t}{2}}} = \frac{|t|^{\frac{1}{2} + \epsilon} e^{(-\operatorname{Re}(\delta) - \pi)t}}{1 - e^{-\pi t}}.$$

This implies that the first integral is analytic for $\operatorname{Re}(\delta) > -\pi$. Similarly, the second integral is analytic for $\operatorname{Re}(\delta) < \pi$. Consequently, both integrals of (5.22) are analytic in δ in $|\delta| < \pi$.

From (5.21) and above discussion, we can say that the expression

$$\begin{aligned}
\phi(\delta) &:= \int_0^\infty \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt + \pi e^{-\frac{i\delta}{2}} \sum_{n=1}^\infty d(n) \log(n) \exp(-2\pi i n e^{-i\delta}) \\
&\quad - i e^{\frac{i\delta}{2}} \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12} - \frac{\log^2(2\pi i e^{-i\delta})}{2} \right) \tag{5.23}
\end{aligned}$$

is an analytic function of δ in $|\delta| < \pi$ except at $\delta = -\frac{\pi}{2}$, because the third term of (5.23) is not analytic at $\delta = -\frac{\pi}{2}$. Hence, $\phi(\delta) + i e^{\frac{i\delta}{2}} \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12} - \frac{\log^2(2\pi i e^{-i\delta})}{2} \right)$ can be expressed as a power series in δ for $\{\delta : |\delta| < \pi\} \setminus \{-\frac{\pi}{2}\}$. Letting $y = 2\pi i(e^{-i\delta} - 1)$ in the asymptotic expansion of $\sum_{n=1}^\infty d(n) \log(n) e^{-ny}$ in (1.15), we find that as $\delta \rightarrow 0$,

$$\begin{aligned}
&\pi e^{-\frac{i\delta}{2}} \sum_{n=1}^\infty d(n) \log(n) \exp(-2\pi i n e^{-i\delta}) \\
&= \pi e^{-\frac{i\delta}{2}} \sum_{n=1}^\infty d(n) \log(n) \exp(-2\pi i n(e^{-i\delta} - 1)) \\
&= \pi e^{-\frac{i\delta}{2}} \left[\frac{1}{2\pi i(e^{-i\delta} - 1)} \left(\log^2(2\pi i(e^{-i\delta} - 1)) + \frac{\pi^2}{6} - \gamma^2 \right) - \frac{1}{2} \log(2\pi) \right. \\
&\quad \left. + \sum_{k=2}^{m-1} \frac{B_{2k}(2\pi i(e^{-i\delta} - 1))^{2k-1}}{k} \left\{ \frac{B_{2k}}{(2k)!} \left(\gamma - \sum_{j=1}^{2k-1} \frac{1}{j} + \log(2\pi) \right) + \frac{2(-1)^k \zeta'(2k)}{(2\pi)^{2k}} \right\} \right. \\
&\quad \left. + \frac{2\pi i(e^{-i\delta} - 1)}{6} \left(\log A - \frac{1}{12} \right) + O((e^{-i\delta} - 1)^{2m-1}) \right] \\
&= \frac{1}{4 \sin\left(\frac{\delta}{2}\right)} \left(\log^2(2\pi\delta) + \frac{\pi^2}{6} - \gamma^2 \right) - \frac{\pi}{2} e^{-\frac{i\delta}{2}} \log(2\pi) \\
&\quad + \sum_{k=1}^{m-1} \pi c_k e^{-\frac{i\delta}{2}} (2\pi i(e^{-i\delta} - 1))^{2k-1} + O((e^{-i\delta} - 1)^{2m-1})
\end{aligned}$$

$$= \frac{1}{4 \sin\left(\frac{\delta}{2}\right)} \left(\frac{\pi^2}{6} - \gamma^2 + \log^2(2\pi\delta) \right) + \sum_{k=0}^{2m-2} d_k \delta^k + O(\delta^{2m-1}), \quad (5.24)$$

as $\delta \rightarrow 0$, where c_k and d_k are effectively computable constants. Consequently, from (5.23) and (5.24), as $\delta \rightarrow 0$, we get

$$\int_0^\infty \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt = \frac{1}{4 \sin\left(\frac{\delta}{2}\right)} \left(\gamma^2 - \frac{\pi^2}{6} - \log^2(2\pi\delta) \right) + \sum_{k=0}^{2m-2} d_k \delta^k + O(\delta^{2m-1}).$$

This completes the proof. \square

6. CONCLUDING REMARKS

The main highlight of this paper was to derive (1.13), that is, an exact transformation for the series $\sum_{n=1}^\infty \frac{\log(n)}{e^{ny} - 1}$, or equivalently, for $\sum_{n=1}^\infty d(n) \log(n) e^{-ny}$, where $\operatorname{Re}(y) > 0$. Such a transformation was missing from the literature up to now. One of the reasons for this could be the lack of availability of the transformation in (1.5) until [23] appeared, which, in turn, resulted from evaluating an integral with a combination of Bessel functions as its associated kernel. This underscores the importance of the applicability of integral transforms in number theory.

In addition, it is to be emphasized that in order to derive (1.13), several new intermediate results, interesting in themselves, had to be derived, for example, the ones given in Theorems 3.3, 3.4, and 4.3. This also shows how important $\psi_1(z)$, and, in general, $\psi_k(z)$, are in number theory, and further corroborates Ishibashi's quote given in the introduction.

The transformation in (1.13) has a nice application in the study of moments of $\zeta(s)$, namely, to obtain the asymptotic expansion of $\int_0^\infty \zeta\left(\frac{1}{2} - it\right) \zeta'\left(\frac{1}{2} + it\right) e^{-\delta t} dt$ as $\delta \rightarrow 0$. However, considering the fact that (1.13) holds for *any* y with $\operatorname{Re}(y) > 0$, we expect further applications of it in the future, in particular, in the theory of modular forms.

A generalization of (1.13), equivalently, of (1.14), may be obtained for the series $\sum_{n=1}^\infty d(n) \log^k(n) e^{-ny}$. This would require differentiating (1.5) k times with respect to a and then letting $a \rightarrow 0$. This would involve Ishibashi's higher analogues of Deninger's $R(z)$ defined in (3.4). Ishibashi [36, Theorem 1] gave a Plana-type formula for $R_k(x)$, $x > 0$, which is easily seen to hold for complex x in the right-half plane $\operatorname{Re}(x) > 0$. This formula involves a polynomial in $\log(t)$ defined by $S_k(t) := \sum_{j=0}^{k-1} a_{k,j} \log^j(t)$, where $a_{k,j}$ are recursively defined by $a_{k,j} = -\sum_{r=0}^{k-2} \binom{k-1}{r} \Gamma^{(k-r-1)}(1) a_{r+1,j}$, $0 \leq j \leq k-2$, with $a_{1,0} = 1$ and $a_{k,k-1} = 1$.

Observe that $S_1(t) = 1$ and $S_2(t) = \gamma + \log(t)$, and so the numerators of the summands of the series on the left-hand sides of (1.6) and (1.14) are $S_1(ny)$ and $S_2(ny)$ respectively. In view of this, we speculate the left-hand side of (1.14) to generalize to $\sum_{n=1}^\infty \frac{S_{k+1}(ny)}{e^{ny} - 1}$ and the right-hand side to involve $\psi_k(z)$.

Since (5.18) can be generalized to

$$\sum_{n=1}^\infty \frac{\log^k(n)}{e^{ny} - 1} = -\frac{1}{2\pi i} \int_{(c)} \Gamma(s) \zeta(s) \zeta^{(k)}(s) y^{-s} ds \quad (c = \operatorname{Re}(s) > 1, \operatorname{Re}(y) > 0),$$

on account of the fact that $\zeta^{(k)}(s) = (-1)^k \sum_{n=1}^\infty \log^k(n) n^{-s}$, once a generalization of (1.14) is obtained, it would possibly give us the complete asymptotic expansion of $\int_0^\infty \zeta\left(\frac{1}{2} - it\right) \zeta^{(k)}\left(\frac{1}{2} + it\right) e^{-\delta t} dt$ as $\delta \rightarrow 0$. Details will appear elsewhere.

Acknowledgements

The authors sincerely thank Christopher Deninger, Nico M. Temme, Rahul Kumar, Shigeru Kanemitsu, Alessandro Languasco for helpful discussions. They also thank Professor Inese Bula from University of

Latvia for sending a copy of [54]. Part of this work was done when the first author was a National Post-doctoral Fellow (NPDF) at IIT Gandhinagar funded by the grant PDF/2021/001224. He is currently an INSPIRE faculty at IISER Kolkata supported by the DST grant DST/INSPIRE/04/2021/002753. The second author's research is funded by the Swarnajayanti Fellowship grant SB/SJF/2021-22/08. The third author is supported by CSIR SPM Fellowship under the grant number SPM-06/1031(0281)/2018-EMR-I. All of the authors sincerely thank the respective funding agencies for their support.

REFERENCES

- [1] A. Apelblat, *Differentiation of the Mittag-Leffler functions with respect to parameters in the Laplace transform approach*, Mathematics, 8 (2020), 657.
- [2] T. Apostol, *Introduction to analytic number theory*, Springer Science+Business Media, New York, 1976.
- [3] F. V. Atkinson, *The mean value of the zeta-function on the critical line*, Q. J. Math. (Oxford) **10** (1939), 122–128.
- [4] S. Banerjee and R. Kumar, *Explicit identities on zeta values over imaginary quadratic field*, submitted for publication, arXiv: 2105.04141, 2021.
- [5] B. C. Berndt, *On the Hurwitz zeta function*, Rocky Mountain J. Math. **2** (1972), 151–157.
- [6] B. C. Berndt, *On Eisenstein series with characters and the values of Dirichlet L-functions*, Acta Arith. **28** No. 3 (1975), 299–320.
- [7] B. C. Berndt, *Modular transformations and generalizations of several formulae of Ramanujan*, Rocky Mountain J. Math. **7** (1977), 147–189.
- [8] B. C. Berndt, *Ramanujan's notebooks, Part I*, Springer-Verlag, New York, 1985.
- [9] B. C. Berndt, *Ramanujan's Notebooks, Part II*, Springer-Verlag, New York, 1989.
- [10] B. C. Berndt and A. Straub, *Ramanujan's formula for $\zeta(2n + 1)$* , Exploring the Riemann zeta function, Eds. H. Montgomery, A. Nikeghbali, and M. Rassias, pp. 13–34, Springer, 2017.
- [11] I. V. Blagouchine, *A theorem for the closed-form evaluation of the first generalized Stieltjes constant at rational arguments and some related summations*, J. Number Theory **148** (2015), 537–592.
- [12] D. M. Bradley, *Series acceleration formulas for Dirichlet series with periodic coefficients*, Ramanujan J. **6** (2002), 331–346.
- [13] Yu. A. Brychkov, O. I. Marichev and N. V. Svischenko, *Handbook of Mellin Transforms*, Advances in Applied Mathematics, CRC Press, Boca Raton, FL, 2019, 586 pp.
- [14] T. Chatterjee and S. S. Khurana, *Shifted Euler constants and a generalization of Euler-Stieltjes constants*, J. Number Theory **204** (2019), 185–210.
- [15] M. W. Coffey, *Series representations for the Stieltjes constants*, Rocky Mountain J. Math. **44** No. 2 (2014), 443–477.
- [16] E. T. Copson, *Theory of Functions of a Complex Variable*, Oxford University Press, Oxford, 1935.
- [17] C. Deninger, *On the analogue of the formula of Chowla and Selberg for real quadratic fields*, J. Reine Angew. Math. **351** (1984), 171–191.
- [18] K. Dilcher, *Generalized Euler constants for arithmetical progressions*, Math. Comp. **59** (1992), 259–282.
- [19] K. Dilcher, *On generalized gamma functions related to the Laurent coefficients of the Riemann zeta function*, Aequationes Math. **48** (1994), 55–85.
- [20] A. Dixit and R. Gupta, *Koshliakov zeta functions I. Modular relations*, Adv. Math. **393** (2021), Paper No. 108093.
- [21] A. Dixit, R. Gupta and R. Kumar, *Extended higher Herglotz functions I. Functional equations*, submitted for publication.
- [22] A. Dixit, R. Gupta, R. Kumar and B. Maji, *Generalized Lambert series, Raabe's cosine transform and a two-parameter generalization of Ramanujan's formula for $\zeta(2m + 1)$* , Nagoya Math. J. **239** (2020), 232–293.
- [23] A. Dixit, A. Kesarwani and R. Kumar, *Explicit transformations of certain Lambert series*, Research in the Mathematical Sciences **9**, 34 (2022).
- [24] A. Dixit and B. Maji, *Generalized Lambert series and arithmetic nature of odd zeta values*, Proc. Royal Soc. Edinburgh, Sect. A: Mathematics, **150** Issue 2 (2020), 741–769.
- [25] M. M. Dzhrbashyan, *Integral Transforms and Representations of Functions in the Complex Domain*, Izdat. "Nauka", Moscow, 1966, 671 pp.
- [26] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Tables of integral transforms. Vol. I*. Based, in part, on notes left by Harry Bateman. McGraw-Hill Book Co., Inc., New York-Toronto-London, 1954.
- [27] J. W. L. Glaisher, *On the product $1^1.2^2.3^3 \dots n^n$* , Mess. Math. (2)**VII** (1877), 43–47.
- [28] J. W. L. Glaisher, *On the constant which occurs in the formula for $1^1.2^2.3^3 \dots n^n$* , Mess. Math. **24** (1894), 1–16.
- [29] S. M. Gonek, *Mean values of the Riemann zeta function and its derivatives*, Invent. Math. **75** No. 1 (1984), 123–141.
- [30] R. Gorenflo, A. A. Kilbas, F. Mainardi and S. V. Rogosin, *Mittag-Leffler Functions, Related Topics and Applications*, Springer Monographs in Mathematics, Springer-Verlag 2014, 443 pp.

- [31] I.S. Gradshteyn and I.M. Ryzhik, eds., *Table of Integrals, Series, and Products*, 7th ed., Academic Press, San Diego, 2007.
- [32] A. Gupta and B. Maji, *On Ramanujan's formula for $\zeta(1/2)$ and $\zeta(2m+1)$* , J. Math. Anal. Appl. **507** (2022), 125738.
- [33] G. H. Hardy and J. E. Littlewood, *Contributions to the Theory of the Riemann Zeta-Function and the Theory of the Distribution of Primes*, Acta Math., **41** (1916), 119–196.
- [34] G. H. Hardy and J. E. Littlewood, *On Lindelöf's hypothesis concerning the Riemann zeta-function*, Proc. Royal Soc. London (A) **103** (1923), 403–412.
- [35] A. E. Ingham, *Mean-value theorems in the theory of the Riemann zeta-function* Proc. London Math. Soc. **27** (1928), 273–300.
- [36] M. Ishibashi, *Laurent coefficients of the zeta function of an indefinite quadratic form*, Acta Arith. **106** No. 1 (2003), 59–71.
- [37] M. Ishibashi and S. Kanemitsu, *Dirichlet series with periodic coefficients*, Res. Math. **35** (1999), 70–88.5
- [38] A. Ivić, *The mean values of the Riemann zeta-function on the critical line*, Analytic Number Theory, Approximation Theory, and Special Functions, 3–68, Springer, New York, 2014.
- [39] E. Jahnke and F. Emde, *Tables of functions with formulae and curves*, 4th ed., Dover Publications, New York, 1945.
- [40] D. S. Jones, *The Theory of Electromagnetism*, MacMillan, New York, 1964.
- [41] S. Kanemitsu, Y. Tanigawa and M. Yoshimoto, *Ramanujan's formula and modular forms* Number Theoretic Methods (Iizuka, 2001), 159–212, Dev. Math., 8, Kluwer Acad. Publ., Dordrecht, 2002.
- [42] K. Katayama, *Ramanujan's formulas for L -functions*, J. Math. Soc. Japan **26** No. 2 (1974), 234–240.
- [43] H. Kinkelin, *Ueber eine mit der Gammafunction verwandte transcendente und deren anwendung auf die integralrechnung*, J. Reine Angew. Math. **57** (1860), 122–138.
- [44] Y. Komori, K. Matsumoto and H. Tsumura, *Barnes multiple zeta-functions, Ramanujan's formula, and relevant series involving hyperbolic functions*, J. Ramanujan Math. Soc. **28** No. 1 (2013), 49–69.
- [45] A. Languasco and L. Righi, *A fast algorithm to compute the Ramanujan-Deninger gamma function and some number-theoretic applications*, Math. Comp. **90** No. 332 (2021), 2899–2921.
- [46] F. Oberhettinger, *Tables of Mellin Transforms*, Springer-Verlag, New York, 1974.
- [47] F. W. J. Olver, *Asymptotics and Special Functions*, AKP Classics, reprint of the 1974 original [Academic Press, New York], CRC Press, 2010.
- [48] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark, eds., *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.
- [49] R. B. Paris and D. Kaminski, *Asymptotics and Mellin-Barnes Integrals*, Encyclopedia of Mathematics and its Applications, 85. Cambridge University Press, Cambridge, 2001.
- [50] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, *Integrals and series*, Vol. 1: Elementary Functions, Gordon and Breach, New York, 1986.
- [51] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, *Integrals and series*, Vol. 3: More Special Functions, Gordon and Breach, New York, 1990.
- [52] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa, New Delhi, 1988.
- [53] S. Ramanujan, *Notebooks* (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957; second ed., 2012.
- [54] E. Riekstins, *Asymptotic expansions for some type of integrals involving logarithms*, Latvian Math. Yearbook **15** (1974), 113–130.
- [55] S. Shirasaka, *On the Laurent coefficients of a class of Dirichlet series*, Result. Math. **42** (2002), 128–138.
- [56] K. Soundararajan, *Moments of the Riemann zeta function*, Ann. Math. **170** No. 2 (2009), 981–993.
- [57] N. M. Temme, *Asymptotic Methods for Integrals*, Series in Analysis, Vol. 6, World Scientific, Singapore, 2015.
- [58] E. C. Titchmarsh, *The Theory of the Riemann Zeta Function*, Clarendon Press, Oxford, 1986.
- [59] C. L. Turnage-Butterbaugh, *Gaps between zeros of Dedekind zeta-functions of quadratic number fields*, J. Math. Anal. Appl. **418** (2014), 100–107.
- [60] A. Voros, *Spectral Functions, Special Functions and the Selberg zeta function*, Commun. Math. Phys. **110** (1987), 439–465.
- [61] A. Wiman, *Über den fundamental satz in der theorie der funcktionen $E_a(x)$* , Acta Math. **29** (1905), 191–201.
- [62] S. Wigert, *Sur la série de Lambert et son application à la théorie des nombres*, Acta Math. **41** (1916), 197–218.
- [63] R. Wong and M. Wyman, *A generalization of Watson's lemma*, Canad. J. Math. **24** No. 2 (1972), 185–208.

DISCIPLINE OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, GANDHINAGAR, PALAJ, GANDHINAGAR 382355, GUJARAT, INDIA

Current address: Department of Mathematics and Statistics, Indian Institute of Science Education and Research Kolkata, Mohanpur, Nadia 741246, West Bengal, India

Email address: soumya.tatan@gmail.com

DISCIPLINE OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GANDHINAGAR, PALAJ, GANDHINAGAR 382355, GUJARAT, INDIA

Email address: adixit@iitgn.ac.in

DISCIPLINE OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GANDHINAGAR, PALAJ, GANDHINAGAR 382355, GUJARAT, INDIA

Email address: shivajee.o@iitgn.ac.in