


(1)

## Principle of deformation of Path:

This idea is related to path independence. We may



imagine that the path  $C_2$  is obtained from  $C_1$  ~~is~~ by continuously moving  $C_1$  (with ends fixed)

until it coincides with  $C_2$ . The integrals along these paths remain unchanged.

Hence, we may conduct a continuous deformation of path of an integral,

keeping ends fixed. So long as the deforming path always contains points at which  $f(z)$  is analytic (inside a simply connected domain) the

integral retains the same value; this is called the principle of deformation of path.

Example : Using the principle we can show:

$$\oint_C (z - z_0)^m dz = \begin{cases} 2\pi i & (m = -1) \\ 0 & (m \neq -1) \end{cases}$$

m integer

for counterclockwise integration around any simple closed path  $C$ , containing  $z_0$

in its interior

In fact, for ~~small~~ <sup>some</sup>  $\epsilon > 0$ ,

the circle,  $|z - z_0| = \epsilon$  can

be continuously deformed

in two steps, into a

path, just indicated, by

first deforming one semicircle

and then the other.

Cauchy's integral theorem ③  
for multiply connected  
domains:

Cauchy's theorem applies  
to multiply connected  
domains. We first explain  
this for a doubly connected  
domain  $D$  with outer boundary  
 $C_1$  and inner boundary  $C_2$ .



If a function  
 $f(z)$  is  
analytic in  
any domain


$D^*$  that contains  $D$  and  
its boundary curves, we  
claim that,  $\oint_{C_1} f(z) dz = \oint_{C_2} f(z) dz$ ,  
both integrals taken  
counterclockwise; here

(4)

full interior of  $C_2$  may not belong to  $D^*$ .

Proof:

By two cuts



$L_1$  and  $L_2$  we split  $D$  into two simply

connected domains  $D_1$  and  $D_2$  in which and on whose boundaries,  $f(z)$  is analytic. By Cauchy's integral theorem, the integral over the entire boundary of  $D_1$  (taken in the sense of arrows of figure) is zero and so is the integral over the boundary of  $D_2$ .

(5)


Hence the sum of these integrals is zero. In this sum the integrals along the cuts  $L_1$  and  $L_2$  cancel, since we integrate along them in opposite directions; we are left with the integral along  $C_1$  (counter clockwise) and along  $C_2$  (clockwise). Hence, reversing the integration along  $C_2$ , we find :

$$\oint_{C_1} f(z) dz - \oint_{C_2} f(z) dz = 0.$$

This completes the proof.

For domains of higher connectivity idea remains the same. Thus, for a triply connected domain, we use three cuts

(6)



Adding integrals as before, the integrals over cuts cancel and the integrals over  $C_1$  (counter-clockwise) and over  $C_2, C_3$  (clockwise) have sum equal to zero.


Hence the integral over  $C_1$  equals the sum of integrals over  $C_2$  and  $C_3$ , all taken counter-clockwise.

Revisit:  $\oint_C (z - z_0)^m dz$ .

## Cauchy's Integral formula: (7)

The formula is used for evaluating integrals of certain type. It has good consequences also: analytic functions have derivatives of all orders.

Theorem: Let  $f(z)$  be analytic in a simply connected domain  $D$ . If  $z_0$  is any point in  $D$  and  $C$  is any simple closed path in  $D$  which encloses  $z_0$ , then



$$\oint_C \frac{f(z)}{z - z_0} dz = 2\pi i \cdot f(z_0)$$

Where integration is taken counter-clockwise.

(8)

We may rewrite the formula:

$$f(z_0) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z-z_0} dz.$$

This is Cauchy's integral formula.

Proof depends on Cauchy's integral theorem, some inequalities and ML-inequality.

Examples:

(1)  $\oint_C \left( \frac{\sin z}{z - \frac{\pi}{2}} \right) dz$

$$= 2\pi i \cdot \sin \frac{\pi}{2} = 2\pi i,$$

for any counterclockwise

contour C enclosing  $\frac{\pi}{2}$ , since  
 $\sin z$  is entire; if C has

$\frac{\pi}{2}$  outside it, value is zero.

(9)

$$\begin{aligned}
 (2) \quad & \oint_C \frac{3z-4}{2z^2-7z+3} dz, \quad C: |z|=2 \text{ counterclockwise} \\
 &= \oint_C \left( \frac{1}{2z-1} + \frac{1}{z-3} \right) dz \\
 &= \oint_C \left( \frac{1}{2z-1} \right) dz + \oint_C \left( \frac{1}{z-3} \right) dz \\
 &= \frac{1}{2} \oint_C \left( \frac{1}{z-\frac{1}{2}} \right) dz + \oint_C \left( \frac{1}{z-3} \right) dz \\
 &= \frac{1}{2} \cdot 2\pi i \cdot 1 + 0 = \pi i.
 \end{aligned}$$


(First integral has  $\frac{1}{2}$  enclosed by  $C$ , hence Cauchy's integral formula applies with  $f(z)=1, \forall z$ .

Second integral has 3 lying outside  $C$ , hence Cauchy's integral theorem applies:  $\frac{1}{z-3}$  is analytic inside the simply connected domain  $D: |z| < \frac{5}{2}$ , in which  $C$  lies.)

(10)

(3) Integrate  $g(z) = \frac{z^2 + 1}{z^2 - 1}$

Clockwise around each of the four 'circles' in



the figure

here.



Note that  $g(z) = \frac{z^2 + 1}{(z-1)(z+1)}$ ,

so  $g(z)$  is analytic everywhere in  $\mathbb{C}$  except at  $z=1$  &  $z=-1$ .

We take care to keep these points in our mind.  
Let us consider each circle separately:

(a) The circle,  $|z-1|=1$  encloses the point  $z_0=1$  (only) where  $g(z)$  is not analytic. Hence, we represent  $g(z)$  as

$$g(z) = \left( \frac{z^2+1}{z+1} \right) \cdot \frac{1}{z-1} = \frac{f(z)}{z-1}$$

Where  $f(z) = \frac{z^2+1}{z+1}$ , is analytic in some simply connected domain, containing the circle  $|z-1|=1$ . Thus

$$\oint_C \left( \frac{z^2+1}{z^2-1} \right) dz = 2\pi i \cdot f(1) = 2\pi i.$$

(b) Answer is same as (a) by principle of deformation of path.

(c) The function  $g(z)$  is same as before; since the 'circle' (simple, closed path) encloses

$z_0 = -1$  (only), we may

express  $g(z)$  as

$$g(z) = \frac{h(z)}{z+1}$$

where  $h(z) = \frac{z^2+1}{z-1}$ .

Thus:  $\oint_C g(z) dz = \oint_C \left( \frac{z^2+1}{z-1} \right) \cdot \frac{1}{z+1} dz$

$$= \oint_C \frac{h(z)}{z+1} dz = 2\pi i \cdot h(-1)$$


$$= 2\pi i (-1) = -2\pi i.$$

(d) Answer is 0, by ---

(4) Integrate counterclockwise:

$$\int_C \frac{\ln(z+1)}{z^2+1}, \quad c: |z-i| = \frac{5}{4},$$

Ans:  $\left( \frac{\ln(i+1)}{i+1} \right) \cdot 2\pi i = \pi \left( \ln \sqrt{2} + \frac{i\pi}{4} \right)$



$$\left| \frac{3-i}{4} \right| = \sqrt{(3)^2 + (-1)^2} = \sqrt{10}$$

$$|3-i| = \sqrt{10} \cdot 4 = \frac{5}{4}$$

$$(3+i) \pi = \arg(3+i)$$


Multiply connected domains can be treated as in discussion relating to Cauchy's integral theorem for such domains:

If  $f(z)$  is analytic on paths  $C_1, C_2$  and in the ring shaped

domain bounded

by them, as

shown in adjoining figure.



If  $z_0$  is any point in the domain  $D$ , then


$$f(z_0) = \frac{1}{2\pi i} \oint_{C_1} \frac{f(z)}{z - z_0} dz + \frac{1}{2\pi i} \oint_{C_2} \frac{f(z)}{z - z_0} dz,$$

where the outer integral over  $C_1$  is taken counterclockwise and the inner clockwise, as indicated in figure.

(14)

Ex: If  $C_1$  is  $|z|=2$ , counterclockwise  
and  $C_2$  is  $|z|=1$  clockwise,  
then

$$\begin{aligned}
 & \oint_{C_1} \frac{e^{z^2}}{z^2(z-1-i)} dz + \oint_{C_2} \frac{e^{z^2}}{z^2(z-1-i)} dz \\
 &= 2\pi i \left[ \frac{e^{z^2}}{z^2} \right]_{z=1+i} \\
 &= 2\pi i \left[ \frac{e^{2i}}{2i} \right] \\
 &= \pi e^{2i}.
 \end{aligned}$$



→ ← X → ←